Changes in the Expression of DNMTs Before and After Treatment with Methotrexate (MTX)/Mercaptopurine (6-MP) in B-Cell ALL Children

Mahmoud Imanei-Avaz¹, Azam Sadat Hashemi², Nasrin Ghasemi³, Seyed Hossein Hekmati Moghaddam⁴, Fatemeh Pourrajab⁵*, Kazem Barzegar⁶, Mahmood Vakili⁷

1. International Campus of Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

2. Hematology & Oncology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

3. Abortion Research Center, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

4. Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

5. Department of Laboratory Sciences, School of Paramedicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

6. English Language Department, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

7. MD, MPH, Associate Professor in Community Medicine, Health Monitoring Research Center, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

*Corresponding authors: Dr Fatemeh Pourrajab, Reproductive Immunology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. E-mail: mina_poorrajab@yahoo.com (F. Pourrajab). ORCID ID: 000-0002-1724-5392

Received: 16 July 2023 Accepted: 18 February 2024

Abstract

Background: DNA methylation is catalyzed by DNA methyltransferases (DNMTs) which are encoded by DNMT1, DNMT3A, and DNMT3B. DNMTs play a major role in the abnormal methylation of tumor suppressors and cancer-related genes. Herein, this study explored the expression profile of DNMTs in pediatric patients with B-cell acute lymphoblastic leukemia (ALL), before and after methotrexate (MTX)/mercaptopurine (6-MP) treatment.

Materials and Methods: This before-after prospective study included 30 matched children in sex and age (20 children with B-cell ALL and 10 healthy children used as a control or calibrator group). The expression profile of DNMTs was assessed at two-time points; at the diagnosis time and after MTX/6-MP treatment in the consolidation-maintenance phase of therapy. Notable, all pediatric patients included in this study continued the therapy without adverse events, except two children who were excluded from the study.

Results: The average age of the patient group was 7.1 ± 1.3 years (in the range of 4-9 years), and the average age of the control group was 8.3 ± 1.7 years (6-10 years). The expression profile of DNMTs in B-cell ALL children was obtained completely different from that in the healthy group. After MTX/6-MP treatment of B-cell ALL children, the expression levels of DNMT1 and 3A were increased (p <0.01 & 0.04, respectively), and the expression level of DNMT3B was decreased (p <0.01), significantly.

Conclusions: In ALL, the expression profile of DNMTs would be changed whereby contribute to abnormal growth and maturation capacity of leukemic stem cells and MTX/6-MP treatment could reverse this profile from a cancerous phenotype to the normal one.

Keywords:	DNA	Methyltransferases	(DNMTs),	Methotrexate	(MTX),	Mercaptopurine	(6-MP)
-----------	-----	--------------------	----------	--------------	--------	----------------	--------

Introduction

Cancer in children becomes more Acute lymphoblastic leukemia (ALL) is the most common cancer diagnosed in children, with certain subsets experiencing greater than 98% cure rate. Most ALL pediatric patients show B-cell immunophenotype (~ 80%) which, in turn, encompasses a broad range of patients with a high cure rate in chemotherapy (1, 2). Today, the standard protocols for ALL chemotherapy start with an induction regimen (~1 month), followed by a consolidation phase (~1-2 months), and finally end with the methotrexate (MTX)/mercaptopurine (6-MP) maintenance therapy which is given until 2 to 3 years from diagnosis. Consolidationmaintenance therapy with anti-metabolites of folate (methotrexate (MTX)) and 6-MP as an effective anticancer drug belongs thiopurine agents with to remarkable success in clinical practice, especially in the treatment of ALL in children. Thiopurines and MTX are widely understood to act as DNA anti-methylating agents (4-6). DNA is methylated at CpG dinucleotides whose clusters (CpG islands) are concentrated in the upstream regions of many anti-cancer genes including tumor suppressors wherein CpG sites are mainly un-methylated and their increased methylation is associated with reduced expression of downstream genes. In cancers, hypermethylation of CpG islands the upstream regions of tumor in suppressors and DNA-repair genes is common, as compared with normal tissues. DNA repair genes are frequently repressed in cancers due to hypermethylation of CpG islands within their promoters. In cancers, loss of expression of genes occurs about 10 times more frequently by hypermethylation of promoter CpG islands than by mutations (7, 8). Methylation of CpG islands is catalyzed by DNA methyltransferases (DNMTs) whose changes in expression have been observed in multiple human cancers, in particular in leukemia. The human genome encodes DNMT1, DNMT3A, and DNMT3B whose activity is responsible for the methylation of DNA and dysregulation results in abnormal methylation of CpG islands (8-10).

Accordingly, MTX/6-MP treatment causes a serious decrease in Sadenosylmethionine (SAM) levels which can directly reduce DNMTs activity and would be closely reflected in the demethylation of CpG islands. Herein, SAM is needed by DNMTs over-expressed in tumors to methylate DNA at special promoters wherein the restriction in SAM levels would result in a reduction in the activity of DNMTs. This could be associated with increased expression of genes inhibiting in tumor cells (e.g. Tumor suppressors (TS), DNA repair, DNMT3A,) (5-8, 11). Additionally, the presence of 6-MP at CpG sites (CpTG), simultaneously hypermethylation enhance may of The above-mentioned oncogenic sites. mechanisms may lead to a reversal of abnormal methylation patterns toward normal ones; anti-methylating functions may help to maintain an active chromatin configuration at CpGs-enriched target loci, for example at the promoter of anti-tumor genes or tumor suppressors which are CpGs rich (12-14).

Hence, to gain biochemical evidence for the hypothesis that DNMTs have an essential role in the cancerous state of ALL and response to chemotherapy, we the assessed expression profile of DNMT1, 3A, & 3B in specimens obtained from B-cell ALL children at diagnosis time and after 3 months of treatment course when pediatric patients were on the consolidation-maintenance phase of MTX/6-MP therapy, and then compared to those in healthy matched children. DNMT expression profile was assessed using Quantitative Real-Time PCR (QRT-PCR) method. The results are expected to lead to a better understanding of the molecular basis of ALL and the epigenetic action of effective chemotherapy. Epigenetic drugs have been of great interest, not only for basic research but also for further understanding of disease pathogenesis and the development of new therapies (10, 15). Therefore, the gene expression levels of DNMT1, 3A, & 3B in specimens obtained from B-cell ALL children were assessed and differences between before and after the treatment and also between healthy and sick people were calculated and compared.

Materials and Methods

79

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

Patients Study groups

This before-after prospective study included 20 pediatric patients with B-cell ALL referred to the Pediatric Oncology Branch of the Children's medical center of Tehran between 2015 and 2016, besides 10 healthy matched children as a control group.

According to similar studies that have reported the mean and standard deviation of DNMT3 gene expression in patients (0.69+0.51) and healthy people (0.35+0.47) with 95% confidence and 80% power for a difference of 0.5 units after the intervention, the number of samples 18 cases were estimated (9).

Therefore, gene expression differences in children with newly diagnosed B-cell acute lymphoblastic leukemia before and after treatment, as well as between healthy and diseased group, were calculated and compared.

The control group consisted of 10 healthy children (8 girls and 2 boys, in the age range of 6-10 years) referred to the central laboratory of Yazd city, and according to the diagnosis of the pediatric oncologist were free of any type of malignancy or disease. The case group included eligible pediatric patients with newly diagnosed Bcell ALL (10 girls and 10 boys, in the age range of 4-9 years) who were previously untreated. Children with any previous chemotherapy or those treated with systemic corticosteroids within 1 month before diagnosis were excluded. Patients with lymphoma syndrome, with massive splenomegaly, or any other malignancy were not eligible for this study. Patients were matched at diagnosis and met the above criteria. Participants were enrolled by convenience sampling, and their blood samples were taken two times. In the case of pediatric patients, the first sampling was at the pre-treatment point when they were newly diagnosed with B-cell ALL and the second one was after 3 months of treatment when patients were receiving dexamethasone (Dex)/vincristine (Vin)/ methotrexate (MTX)/ mercaptopurine (6-MP) chemotherapy drugs for the maintenance therapy, according to the standard regimen protocol invested in the center (2-4, 16).

Sampling and Diagnosis

The research was carried out after obtaining permission of Committee of Ethics in Human Research with code: (IR.SSU.MEDICINE.REC.1394.198-201.

All admitted B-cell ALL children met World Health Organization (WHO) criteria and guidelines for B-cell ALL and were diagnosed and introduced by a pediatric leukemia specialist. About 5 mL of peripheral blood was taken from both pediatric groups, after obtaining written informed consent from all participants' parents.

RNA Isolation

White blood cells (WBC) were isolated by sedimentation Ficoll-Hypaque on gradients. Total RNA was participated and isolated with a Ribo EX kit (Geneall, Daejeon, Korea), according to the manufacturer's instructions. The quantity and purity of RNA content were evaluated (Thermo Fisher using Nanodrop-2000 Scientific. Waltham, MA, USA). Absorbance ratios and concentrations were determined as indicators of sample yield. The integrity and quality of total RNA were controlled with the 1% agarose gel electrophoresis. eliminate DNA To contamination, the isolated RNA solution was treated with DNase (DNA-free, Ambion, Austin-USA) (Figure 1). Gene expression of DNMT1, 3A, and 3B was then assessed by QRT-PCR and analyzed by SPSS15.

Quantitative Real-Time PCR (QRT-PCR)

First-strand cDNA was synthesized from 2 μ g RNA in a 20 μ L reaction mixture containing random hexamers as primers using a cDNA synthesis kit (MBI Fermentas, Ontario–Canada Revert Aid, First Strand cDNA Synthesis Kit). The relative expression of target genes was

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

DOI: 10.18502/ijpho.v14i2.15267

This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use and redistribution provided that the original author and source are credited.

performed by Step One Plus Real-time PCR (Applied Biosystems) in triplicate; primer specificity was confirmed by melting curves (Figure 2). QRT-PCR was done using HOT FIREPol® EvaGreen qPCR Mix (Solis BioDyne, Tartu, Estonia) according to the SYBR Green method. Briefly, QRT-PCR was carried out in 20 µL PCR buffer (10 mM Tris- HCl (pH 9.5), 50 mM KCL, % 0.1 Triton X), 0.2 mM dNTP mix, 1.5 mM MgCl2, 10 pmol of each primer and 2U of Taq-DNA polymerase. The PCR cycle was 94 °C for 30s, 60 °C for 30s and 72 °C for 60s. The primers and their characteristics are described in Table I. To check the primers and cDNA synthesis, the expression of **GAPDH** (Glyceraldehyde Phosphate Dehydrogenase) and **DNMTs** were analyzed by melting curve, and QRT-PCR products were run on a 1% agarose gel stained with EtBr (Figure 1). Relative fold changes in the expression levels of genes were calculated according to the $2^{-\Delta\Delta Ct}$ formula (below), after normalization with GAPDH as a reference gene and then were analyzed by One-way ANOVA.

Statistical Analysis

All the PCR's were carried out in triplicate and the signal intensities were correlated almost linearly with the amount of mRNA The expression levels dilutions. of **DNMTs** normalized with were glyceraldehyde phosphate dehydrogenase (GAPDH) and then the gene expression differences before and after the treatment and also between healthy and sick people were calculated and compared with a t-Test, using SPSS software version 21 (SPSS 21, IBM Corp., Armonk, NY, USA). Differences considered were significant at p-values < 0.05.

Results

Patient Characteristics

The study recruited a total of 30 eligible nearly matched children including 18 girls (10 in the patient group and 8 in the *Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93* healthy group) and 12 boys (10 in the patient group and 2 in the healthy group). However, two of ALL children were excluded from the study because they were not able to continue the treatment and declined participation in the study. All admitted B-cell ALL children met World Health Organization (WHO) criteria and guidelines for B-cell ALL and were diagnosed and introduced by a pediatric leukemia specialist. The mean age of the patient group was 7.1±1.3 years (a range of 4-9 years), and the mean age of the control group was 8.3±1.7 years (a range of 6-10 years) (Table II). Newly diagnosed children with B-cell ALL were sampled before starting the treatment and after 3 months when they were on consolidationmaintenance therapy and receiving dexamethasone (Dex)/vincristine (Vin)/MTX/6-MP chemotherapy drugs, according to the standard regimen protocol invested in the children's medical center (2-4, 16). Except for two children who were excluded from the study, all pediatric patients who were included in this study showed no adverse effects and were responding consolidationto the maintenance regimen. Also, there was no significant difference between the patients showing remission. Herein, in the experiment was divided into two parts: first to investigate the expression profile of DNMT1, 3A, and 3B in B-cell ALL children at diagnosis time before receiving any treatment and compare it with that in healthy children; second, to assess possible changes that occurred in the expression profile of DNMT1, DNMT3A, and DNMT3B in pediatric patients, when they MTX/6-MP consolidationwere on maintenance therapy.

The higher expression level of DNMT1 in B-cell ALL children

This study was conducted on children with newly-diagnosed B-cell ALL sampled before starting the therapy and then after 3 months when they were on consolidation-

maintenance therapy and receiving dexamethasone (Dex)/vincristine (Vin)/MTX/6-MP chemotherapy drugs, according to the standard regimen protocol invested in the children's medical center. Blood samples were taken from 20 children newly diagnosed with B-cell ALL before starting the chemotherapy, and also from 10 healthy matched children, as controls. Total mRNAs were isolated from their WBCs and then the expression levels of DNMTs were assessed as described previously. Figure 1 is a representative pattern of QRT-PCR products obtained under non-saturating conditions from samples. For a better understanding of differences in the expression profile of DNMTs in B-cell ALL children, QRT-PCR data were analyzed with appropriate methods whereby the expression of the healthy group was considered as a calibrator and the expression of GAPDH internal reference. After as an normalization with GAPDH, the fold expression level of DNMT1 in the healthy group was 1.03±0.3, which increased to 1.98±0.09 in the patient group (with P~0.02) (Figure 3 & Table III).

The lower expression level of DNMT3A in B-cell ALL children

The human DNMT3A together with DNMT3B have been associated with de novo methylation of CpG dinucleotides whereby they are establishing genomic imprints. Analysis of DNA methylation patterns in normal and tumor cells has revealed that many human tumor cells have hypermethylation and inactivation of tumor suppressor genes such as p16 or p53 (9, 11). Hence, this study analyzed the fold expression level of DNMT3A in B-cell ALL children and compared it with that in healthy the normal group. After normalization with GAPDH, the fold expression level of DNMT3A in the healthy control group was 2.72±0.22 which was 1.21±0.6 in B-cell ALL children (with P~0.04) (Figure 3 & Table III).

The higher expression level of DNMT3B in B-cell ALL children

Finally, the fold expression level of DNMT3B in blood cells from B-ALL children was analyzed and compared with the healthy group. The relative expression level of DNMT3B in the healthy control group was 1.26 ± 0.62 which was 3.27 ± 0.3 in B-cell ALL children (with P~0.03) (Figure 3 & Table III). Interestingly, consistent with DNMT1 and unlike the expression level DNMT3A, of DNMT3B was significantly higher in pediatric patients than in healthy children. Besides, consistent with DNMT1 and unlike DNMT3A, the expression level of DNMT3B was significantly higher in B-ALL children than in the healthy group

In the next step, the study aimed to assess changes that appeared in the expression profile of DNMTs, after B-ALL children received MTX/6-MP therapy.

In the next step, the study aimed to evaluate the changes that appeared in the expression level of DNMTs in B-ALL children after receiving MTX/6-MP treatment.

Decreased expression of DNMT1 after MTX/6-MP treatment

Previous studies on the epigenome and drug-targeted mechanisms in tumor cells have emphasized the reversibility of promoter hypermethylation of cancerrelated genes. They proposed that hypermethylation of genomic sites can be reversed by treatment with chemotherapy agents and emphasized the role of DNMTs as optimal targets for what is now known as epigenetic therapy (10, 15).

In this part of the study as mentioned earlier, 20 matched B-cell ALL children (10 girls and 10 boys, with an age range of 4-9 years), were included while making sure that the correct types and doses of drugs were given. Except for two children who were excluded from the study, all Bcell ALL children (10 girls and 8 boys) continued participation in this study, showed no adverse effects, responded to

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

DOI: 10.18502/ijpho.v14i2.15267

This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use and redistribution provided that the original author and source are credited.

the chemotherapy, and went on treatment with MTX/6-MP in the consolidationmaintenance phase (according to the protocol regimen in the children's medical center). Thus, it was worthwhile to investigate the expression profile of **DNMTs** these patients. in In the consolidation-maintenance phase of treatment with MTX/6-MP, the expression of DNMT1 in blood cells from B-cell ALL children showed significantly lowered levels when compared to that obtained at diagnosis time before treatment. The fold change expression of DNMT1 decreased from 1.98±0.09 (before treatment) to 1.58 ± 0.12 (after treatment) $(P \sim 0.01)$ (Figure 4 & Table IV).

Increased expression of DNMT3A after MTX/6-MP treatment

In contrast to DNMT1, DNMT3A was upregulated in blood cells from B-cell ALL

children after MTX/6-MP treatment. Then, the fold change expression of DNMT3A in B-cell ALL children after treatment with MTX/6-MP increased from 1.21 ± 0.6 to 2.3 ± 0.5 , with P~0.04, (Figure 4 & Table IV).

Reduced expression of DNMT3B after MTX/6-MP treatment

the MTX/6-MP Similar to DNMT1, treatment lowered significantly the expression of DNMT3B in B-cell ALL children where the fold change expression reached its level in normal cells (Figure 4 & Table IV). Before treatment, the fold change expression of DNMT3B in B-cell ALL children was 3.27±0.62 which was reduced to 1.7±0.08 after treatment with MTX/6-MP (P~0.02) (Figure 4 & Table IV), and thus got close to the fold change expression of normal cells (1.26±0.62).

Table I: Characteristics and sequences of DNMT1, 3A, & 3B and GAPDH primers used in this study.

	1 5 7 7	1	2	
Gene Name	Forward and Revers primers 5` to 3`	NCBI AC. NO.	cDNA Size	
DNMTI	F:CACCAGGCAAACCACCATCAC	NM_001318731.1	168 bp	
	R:AGCGGTCTAGCAACTCGTTCTC			
DNMT3A	F: CTCCATCGTCAACCCTGCTC	NM_022552.4	200 bp	
	R: TCATCACAGGGTTGGACTCG			
DNMT3B	F: TGCTCTGGAGAAAGCTAGGGT	NM_006892.3	198 bp	
	R: TCATCACAGGGTTGGACTCG			
GAPDH	F: GAGCCACATCGCTCAGACAC	NM_001289746.1	150 bp	
	R:CATGTAGTTGAGGTCAATGAAGG			

Table II: General characteristics of the healthy group and B-cell ALL children included in this study.

Characteristics	Control group	B-cell ALL group*	
Age (years), mean <u>+</u> SD	8.3 <u>+</u> 1.7	Before treatment	After treatment
		7.1 <u>+</u> 1.3	7.5 <u>+</u> 1.4
Females	8 (80%)	10 (50%)	10 (50%)
Males	2 (20%)	10 (50%)	8 (50%)
Males	2 (20%)	10 (50%)	8 (50%)

* B-cell acute lymphoblastic leukemia (B-cell ALL).

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

Table III. The fold change expression of *DNMT1/GAPDH*, *DNMT3A/GAPDH* and *DNMT3B/GAPDH*, in B-cell ALL children and in healthy normal group (as a calibrator), are presented here, after normalization to *GAPDH*.

Gene Name	Control group	Patient group	P-values	
DNMT1, mean <u>+</u> SD	1.03±0.16	1.98±0.09	0.02	
<i>DNMT3A</i> , mean <u>+</u> SD	2.72±0.22	1.21±0.6	0.04	
<i>DNMT3B</i> , mean <u>+</u> SD	1.26±0.62	3.27±0.3	0.03	

SD; Standard deviation, GAPDH; glyceraldehyde phosphate dehydrogenase, DNMT; DNA methyltransferase). The fold change expression of each gene in the patient group and the healthy group was analyzed and differences between the groups were compared with t-Test, P < 0.05 was considered significant

Table IV: Comparison of the fold change expressions of DNMTs in B-cell ALL children before and after the treatment. DNMT1/GAPDH, DNMT3A/GAPDH and DNMT3B/GAPDH, in B-cell ALL children before and after treatment with Methotrexate (MTX)/Mercaptopurine (6-MP), and in healthy normal group (as a calibrator), are presented here, after normalization to GAPDH.

Gene Name	Control healthy group	B-ALL group, Before treatment	B-ALL group, After treatment	P-values
<i>DNMT1</i> , mean <u>+</u> SD	1.03±0.16	1.98±0.09	1.58±0.12	0.01
DNMT3A, mean <u>+</u> SD	2.72±0.22	1.21±0.6	2.3±0.5	0.04
<i>DNMT3B</i> , mean <u>+</u> SD	1.26±0.62	3.27±0.36	1.7±0.08	0.02

SD; Standard deviation, GAPDH; glyceraldehyde phosphate dehydrogenase, DNMT; DNA methyltransferase). The fold change expression of each gene in the patient group, before and after the treatment was analyzed and compared, by using paired t-Test, P < 0.05 was considered significant.

Table V: Postulated pathways and related genes are disrupted by promoter hypermethylation and are associated with gene-silencing in cancers.

Pathway	Genes
Cell-Cycle Control	<i>Rb</i> , <i>p16</i> , <i>p15</i> , <i>p14</i> , <i>p73</i>
DNA Repair	MLH1, O6-MGMT, GST-Pi, BRCA1*
Growth Factor Response Inhibition	ER, RAR-beta, SOCS-1
Inducing Apoptosis	DAP kinase, caspase 8, TMS-1

* *Genes abbreviations*: DNA mismatch repair protein Mlh1 or MutL protein homolog 1 (*MLH1*), O6-alkylguanine DNA alkyltransferase (*O6-MGMT*), Glutathione S-transferases pi (*GST-Pi*) Breast cancer type 1 susceptibility protein (*BRCA1*), Suppressor of cytokine signaling 1 (*SOCS1*), and the inhibitor of STAT-induced and Janus kinase–signal (JAK–STAT) pathway (9, 11, 23-25).

84

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93



В



Figure 1. Representative gel patterns of QRT-PCR products of GAPDH, DNMT1, 3A, & 3B obtained under non-saturating conditions, run on a 1% agarose gel and stained with EtBr. A: QRT-PCR product of GAPDH gene (left to right: patients GAPDH, 150 bp, and DNA marker (ladder 50 bp)); B (left to right): The first well is DNA marker (ladder 50 bp). RT-PCR product of the DNMT1 gene (patients, 168 bp, and negative control (RT-)), RT-PCR product of the DNMT3A gene (patients, 200 bp, negative control (RT-)), RT-PCR product of the DNMT3B gene (patients, 198 bp, and negative control (RT-)).

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93



Figure 2. Melting curves of PCR products after cDNA synthesis of GAPDH and DNMT1, 3A and 3B transcripts.



Figure 3. The fold change expression of *DNMT1*, *3A* and *3B* in B-cell ALL children and in normal healthy group (as a control), after normalization to *GAPDH* which was considered as a reference gene and is expressed constantly in every cell type. The fold change expression of each gene in the patient's group was compared with the same gene in the healthy group, by using t-Test. *; P < 0.01 was considered significant. **; P < 0.02 was considered significant. ***; P < 0.03 was considered significant. ***; P < 0.04 was considered significant. GAPDH; glyceraldehyde phosphate dehydrogenase, DNMT; DNA methyltransferase.

86

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93



Figure 4. Comparison of the fold change expression of DNMTs, in B-cell ALL children before and after treatment with Methotrexate (MTX)/Mercaptopurine (6-MP) and in healthy normal group (as a calibrator), after normalization to *GAPDH* which was considered as a reference gene and is expressed constantly in every cell type. The fold change expression of each gene in the patient's group was compared with the same gene in the healthy group, by using t-Test. Also, the fold change expression of each gene in the patient group, before and after the treatment was analyzed and compared, by using t-Test. *; P < 0.01 was considered significant. **; P < 0.02 was considered significant. ***; P < 0.03 was considered significant. ****; P < 0.04 was considered significant. GAPDH; glyceraldehyde phosphate dehydrogenase, DNMT; DNA methyltransferase.

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

Changes in the Expression of DNMTs Before and After Treatment with Methotrexate (MTX)/Mercaptopurine (6-MP) in B-Cell ALL Children



Figure 5. A proposed mechanism by 6-MP/MTX to reactivate silenced genes/inactivate protooncogenic sites is illustrated here. MTX as an analog of folate and 6-MP as an analog of purines cause reducing levels of S-adenosylmethionine (SAM) in the cells. SAM is essentially needed to maintain hypermethylation of specific CpGs by tumor-over-expressed DNMTs, as well as, thiopurine methyltransferase (TPMT) to metabolize 6-MP. TPMT metabolizes 6-MP and like as DNMTs, transfers methyl groups from SAM and generates S-adenosylhomocysteine (SAH). MTX/6-MP treatment causes a serious decrease in SAM levels by which can directly reduce DNMTs activity and would be closely reflected in the demethylation of CpG islands. Herein, SAM is needed by DNMTs over-expressed in tumors to methylate DNA at special promoters wherein the restriction in SAM levels would result in a reduction in the activity of DNMTs and could be associated with increased expression of genes inhibiting in tumor cells (e.g. TS, DNA repair, DNMT3A,...). Additionally, presence of 6-MP at CpG sites (CpTG), simultaneously may enhance inactivation of proto-oncogenic sites. The extent of MTX/6-MP effects is through DNA incorporation of 6-MP and also through reducing concentrations of SAM which is influenced by MTX concentration. CpG, Cytosine-guanine dinucleotide sequence: DNMTs, DNA methyltransferases; SAM, S-adenosylmethionine; 6-MP, 6-Mercaptopurine; MTX, Methotrexate; TPMT, Thiopurine methyltransferase; TG, thioguanine nucleotide; TS, Tumor suppressors.

Discussion

Epigenetic enzymes DNMTs play a major role in promoter methylation and silencing of tumor suppressor genes in human cancer cells. Even more, aberrant methylations by these genes facilitate gene mutation, since the deamination of 5methylcytosine leads to the formation of thymine which is not repaired by glycosylases (9, 17). Tumor suppressors and DNA repair genes have frequent expression in proliferating cells wherein they are required for genome stability;

impairment in their role can lead to the pathogenesis of leukemic stem cells (7-9). Data from ALL patients indicate that there is a significant reduction in the expression of genes involved in DNA repair and that these are responsible for genome stability including tumor suppressor and cell cyclerelated genes. These changes are clear in the lymphocytes of ALL patients and are associated with an increased level of mutations in DNA and induction of transformation during the cell cycle (3, 8, 15).

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

As remarkable instance, CpG а methylation of p15 and p16 promoters has frequently occurred in acute leukemia. Accordingly, about 93% of AML patients, or 61-94% of all leukemia subtypes display methylation of tumor suppressor promoters (18-20). These highly expressed genes have CpG-rich promoters (high-CpG-density class (HCP)) which are usually un-methylated in normal cells, since these genes are needed to control the cell cycle and growth (8, 20). Studies have identified methylated CpG sites in the promoter of these genes which are frequently inactivated in cancers and also silenced in acute leukemia (7, 9, 11). Methylation of CpG islands is catalyzed by DNMT1, 3A, and 3B whose changes in expression are observed in multiple human cancers, in particular in leukemia (9, 10, 21). DNMT1 is a key methyltransferase in the maintenance of genome methylation, whereas DNMT3A, together with DNMT3B, has an essential role in de novo methylation (7, 11). Additionally, studies have indicated that the promoter regions of DNMTs are also enriched in CpG dinucleotides, the best substrates for DNA methyltransferase activity (13,14). However, changes in DNMTs expression can be observed in a cancer state or even after the chemotherapy phase wherein it could be accounted as an effect associated with an anti-cancer mechanism of drugs (5, 6, 12).

Accordingly, CpG methylation is reversible by treatment with antimethylating agents which are regarded as optimal targets in epigenetic therapy. It is apparent from experimental settings that the reactivation of the types of genes epigenetically silenced in cancer would have a profound anti-tumor effect (22-25). **De-methylation** and reactivation of silenced tumor suppressor genes (e.g. CDKN2A) possible is through chemotherapy by anti-methylating agents (15, 26). DNA anti-methylating agents

exhibit remarkable success in clinical practice especially effective in the treatment of ALL. Whether the underlying mechanism leads to global or selective demethylation is not clearly defined yet (3, 5, 6). In this regard, as anti-methylating agents that negatively affect tumor growth, MTX/6-MP therapy is hypothesized to act through a mechanism directly affecting DNMTs activity followed indirectly by changes in the expression of tumor-related genes (3, 5, 6, 12). Note that, MTX/6-MP treatment seriously reduces the level of Sadenosylmethionine (SAM) in the cells, which is essentially needed by DNMTs to catalyze the addition of methyl groups to CpGs sites and is thereby accounted as a mechanism by these drugs (5, 6, 12). Additionally, the level of SAM is also consumed by another enzyme named thiopurine methyltransferase (TPMT) which metabolizes 6-MP and like DNMTs needs SAM to transfer methyl groups and Me-6Mp and generate Sadenosylhomocysteine (SAH) (3, 5, 6). Accordingly, 6-MP can simultaneously be incorporated into CpG dinucleotides and create CpTG sites; this subsequently causes inhibition of DNMT activity and hypermethylation of CpG islands present at the regions of oncogenic properties, identified frequently hypermethylated, and inactivated in cancers and also silenced in acute leukemia (Table V, Figure 5) (6, 8, 12).

However, despite their long history of use, the exact epigenetic mechanism of the MTX/6-MP regimen remains to be clearly defined which certainly improves our understanding of the pharmacology property and gives rise to better chemotherapy use (5, 6, 12).

Thus, at the first stage of this experiment, the expression profile of DNMTs in blood cells from B-cell ALL children was determined and compared to those in healthy children who were used as a control or calibrator group. As expected,

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

the expression profile of DNMTs in B-cell ALL children was obtained completely different from that in healthy children. In B-cell ALL children, the expression levels of DNMT1 and DNMT3B appeared upregulated and had significantly higher levels compared to the healthy group. Previous studies have also reported that up-regulation of these **DNMTs** is associated with tumor progression (20-22). However, the opposite phenotype was observed for DNMT3A which is expected to participate in differentiating events; also, its down-regulation or loss of function has been observed in tumors of breast, prostate, and colorectal lung. carcinomas (11, 20, 21). The second stage of the experiment was to investigate the effect of MTX/6-MP regimen therapy on the expression profile of DNMTs. Then, the expression of DNMT1, 3A, and 3B assessed at the consolidationwas maintenance phase of therapy with MTX/6-MP. The expressions of DNMT1 and 3B appeared coordinately downregulated, whereas the opposite result was observed for DNMT3A expression. It appeared that the profile was changing from a cancerous phenotype toward the normal one and their expressions were reaching those levels in normal cells. It is notable that all pediatric patients included in this study continued the consolidationmaintenance therapy while showing no adverse effects, except two children who were excluded from the study (Figures 3 & 4). Hence, it is concluded that the MTX/6-MP regimen therapy may act through a mechanism that affects the expression and activity of DNMTs (9, 13, 14, 20). It is important to note that DNMT1 knockdown is understood to cause the induction of genes involved in cell cycle arrest and stress response, whose suppression may have important therapeutic implications. It has been also observed that DNMT3B depletion in human cancer cells induces apoptosis, but not in normal cells. DNMT3B activity seems to be associated

with tumor suppressor genes and their promoter silencing (Table V) (22-24). Moreover, according to data, DNMT3B and DNMT1 may act highly cooperative which is needed by tumor cells to methylate anti-cancer and cell-cyclerelated promoters (20-23). DNMT3B and have been detected highly DNMT1 enriched at hypermethylated loci of these genes with high and intermediate CpGdensity promoters (HCPs & ICPs) (8, 9, 20). Regarding data, one of the optimal targets in tumor cells to reactivate silenced genes and lowering DNA methylation in the promoters seems to be disruption of DNMT1 and DNMTB, whereas DNMT3A has appeared with differentiating properties. DNMT3A has been found in pre-centromeric and telomeric regions and has been supposed to be associated with the methylation of these regions; however, it is essentially needed to contract tumor progression (5, 6, 8). The changes identified in the expression profile of DNMTs from ALL pediatric patients in this study may be used as an indirect mechanism by MTX/6-MP drugs used in cancer treatments.

Conclusion

MTX/6-MP regimen in the consolidationmaintenance phase is an effective therapy with anticancer remarkable success in B-cell ALL remission: however, the accurate mechanism of MTX/6-MP remains to be made clear. Regarding previous studies, the extent of MTX/6-MP effects could be through DNA methylation events and tumor-expressed activity of DNMTs, whereby demethylation of CpG islands is selectively induced at promoters of specific genes (2, 4, 25).

It was observed in the present study that the expression profile of DNMTs in B-cell ALL children was completely different from that in the healthy group and that their expressions changed from the cancerous phenotype toward the normal one, after the MTX/6-MP regimen therapy.

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

DOI: 10.18502/ijpho.v14i2.15267

Somehow, the results of this study are expected to lead to a better understanding of the molecular basis of ALL and the epigenetic action of effective chemotherapy. Epigenetic effects of MTX/6-MP drugs have been of great interest, not only for basic research but also for new therapies. However, the results of this work are not the only mechanism by MTX/6-MP; the confirmation of them requires more in vivo studies.

Ethical Consideration

The study protocol was approved by the Institutional Review Board and also by the Ethics Committee at Shahid Sadoughi University of Medical Sciences. The trial registration in the Institutional Review Board and the Medical Ethics Committee was done on 2015-11-25, with registration NO. IR.SSU.MEDICINE.REC.1394.298 (The vice-chancellor of research, at the School of Medicine, Shahid Sadoughi University of Medical Sciences: E-mail: medschool@ssu.ac.ir).

Author's contiributions

Seyed Hossein Hekmati Moghaddam, Azam Sadat Hashemi, & Nasrin Ghasemi: supervising and advising the research and editing the manuscript text;

Mahmoud Imanei: collecting data, doing experiments, and analyzing data;

Fatemeh Pourrajab: supervising the research and writing the main manuscript text.

Mahmood Vakili, and Kazem Barzegar: writing the main manuscript text.

Conflicts of interests

The authors declare no competing interests in this work.

Acknowledgements

The authors thank Ms. Tahereh Ahmadi Afzadi, Zahra Kinzad and Kolan Rahmani

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

for their kind help and all parents of patients who were referred to the Pediatric Oncology Branch of Children's Medical Center of Tehran and participated at the study. Also, we give thanks to the professionals from the Department of Pediatrics Oncology and Hematology center, Shahid Sadoughi Hospital, Shahid Sadoughi University of Medical Sciences, Yazd, Iran, in particular Dr. Ali Dehghani Firoozabadi for their kind supports.

Funding

The authors have no funding source to declare.

References

- Cooper SL, Brown PA. Treatment of Pediatric Acute Lymphoblastic Leukemia. Pediatr Clin North Am 2015; 62(1): 61-73.
- Schmiegelow K, Nielsen SN, Frandsen TL, Nersting J. Mercaptopurine/Methotrexate Maintenance Therapy of Childhood Acute Lymphoblastic Leukemia: Clinical Facts and Fiction. J Pediatr Hematol Oncol 2014; 36: 503-517.
- Coulthard SA, McGarrity S, Sahota K, Berry P, Redfern CP. Three faces of mercaptopurine cytotoxicity in vitro: methylation, nucleotide homeostasis, and deoxythioguanosine in DNA. Drug Metab Dispos 2018; 46(8): 1191-1199.
- 4. Tolar J, Bostrom BC, La ML, Sather HN. Intravenous 6-Mercaptopurine Decreases Salvage After Relapse in Childhood Acute Lymphoblastic Leukemia: А Report from the Children's Cancer Group Study CCG 1922. Pediatr Blood Cancer 2005; 45: 5-9.
- Yuan B, Zhang J, Wang H, Xiong L, Cai Q, Wang T, et al. 6-Thioguanine Reactivates Epigenetically Silenced Genes in Acute Lymphoblastic Leukemia Cells by Facilitating Proteasome-mediated Degradation of 91

DNMT1. Cancer Res 2011; 71(5): 1904-1911.

- Hogarth LA, Redfern CPF, Teodoridis JM, Hall AG, Anderson H, Case MC, et al. The effect of thiopurine drugs on DNA methylation in relation to TPMT expression. Biochem Pharmacol 2008; 76: 1024-1035.
- Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. Cancer genome landscapes. Sci 2013; 339 (6127): 1546-1558.
- Boultwood J, Wainscoat JS. Gene silencing by DNA methylation in haematological malignancies. Br J Haematol 2007; 138(1): 3-11.
- 9. Sayin DB, Kurekc E, Karabulut HG, Ezer U, Bokesoy I. DNA methyltransferase expression differs with proliferation in childhood acute lymphoblastic leukemia. Mol Biol Rep 2010; 37: 2471-2476.
- 10. Szyf M. Epigenetics, DNA Methylation, and Chromatin Modifying Drugs. Annu Rev Pharmacol Toxicol 2009; 49: 243–263.
- 11. Kim MS, Kim YR, Yoo NJ, Lee SH. Mutational analysis of DNMT3A gene in acute leukemia and common solid cancers. APMIS 2012; 121: 85-94.
- 12. Wang H, Wang Y. 6-Thioguanine Perturbs Cytosine Methylation at the CpG Dinucleotide Site by DNA Methyltransferases in Vitro and Acts as a DNA Demethylating Agent in Vivo. Biochem 2009; 48 (10): 2290-2299.
- 13. Lee SJ, Jeon HS, Jang JS, Park SH, Lee GY, Lee BH, et al. DNMT3B polymorphisms and risk of primary lung cancer. Carcinogenesis 2005; 26(2): 403-409.
- 14. Zampieri M, Passananti C, Calabrese P, Perilli M, Corbi N, Cave F, et al. Parp1 Localizes within the Dnmt1 Promoter and Protects Its Unmethylated State by Its Enzymatic

Activity. Plos One 2009; 4(3): e4717-e4721.

- Dunwell TM, Hesson LB, Pavlova T, Zabarovska V, Kashuba V, Catchpoole D, et al. Epigenetic analysis of childhood acute lymphoblastic leukemia. Epigenetics 2009; 4(3): 185-193.
- 16. Sparrow A, Geelhoed G. Prednisolone versus dexamethasone in croup: a randomised equivalence trial. Arch Dis Child 2006; 91(7):580-583.
- 17. Foulks JM, Parnell KM, Nix RN, Chau S, Swierczek K, Saunders M, et al. Epigenetic Drug Discovery: Targeting DNA Methyl transferases. J Biomol Screen 2012; 17(1): 2-17.
- Yang Y, Takeuchi S, Hofmann WK, Ikezoe T, van Dongen JJ, Szczepański T, et al. Aberrant methylation in promoter-associated CpG islands of multiple genes in acute lymphoblastic leukemia. Leu res 2006; 30(1):98-102.
- 19. Stanczyk M, Sliwinski T, Trelinska J, Cuchra M, Markiewicz L, Dziki L, et al. Role of base-excision repair in the treatment of childhood acute lymphoblastic leukaemia with 6mercaptopurine and high doses of methotrexate. Mutat Res 2012; 741(1-2): 13-21.
- 20. Jin B, Ernst J, Tiedemann RL, Xu H, Sureshchandra S, Kellis M, et al. Linking DNA Methyltransferases to Epigenetic Marks and Nucleosome Structure Genome-wide in Human Tumor Cells. Cell Rep 2012; 2: 1411-1424.
- 21. Rostamian T, Pourrajab F, Hekmatimoghaddam S. The effect of 6-thioguanine on proliferation, viability and expression of the genes DNMT 3A, DNMT 3B and HDAC3 in Lymphoid Cancer Cell Line Nalm6. Iran J Pediatr Hematol Oncol 2020; 10(1): 28-37.
- 22. Merry CR, Forrest ME, Sabers JN, Beard L, Gao XH, Hatzoglou M, et al. DNMT1-associated long non-coding

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93

92

RNAs regulate global gene expression and DNA methylation in colon cancer. Hum mol genet 2015; 24(21): 6240-6253.

- Rajabi H, Tagde A, Alam M, Bouillez A, Pitroda S, Suzuki Y, etal. DNA methylation by DNMT1 and DNMT3b methyltransferases is driven by the MUC1-C oncoprotein in human carcinoma cells. Oncogene 2016; 35(50): 6439-6445.
- Eglen RM, Reisine T. Screening for compounds that Modulate epigenetic regulation of the transcriptome: An overview. J Biomol Screen 2011; 16(10): 1137-1152.
- 25. Sanaei M, Kavoosi F, Moezzi MA. Effect of 5'-fluoro-2'-deoxycytidine and sodium butyrate on the genes of the intrinsic apoptotic pathway, p21, p53, cell viability, and apoptosis in human hepatocellular carcinoma cell lines. Iran J Pediatr Hematol Oncol 2021; 11(4): 216-230.
- 26. Bhatla T, Wang J, Morrison DJ, Raetz EA, Burke MJ, Brown P, et al. Epigenetic reprogramming reverses the relapse-specific gene expression signature and restores chemosensitivity in childhood B-lymphoblastic leukemia. Blood 2012; 119(22): 5201-5210.

Iran J Ped Hematol Oncol. 2024, Vol 14, No 2, 78-93