

Socioeconomic Status and Childhood Leukemia

Dehghani Kh MSc¹, Poormovahed Z MSc¹, Dehghani H MSc¹

1- Nursing and Midwifery College, Shahid Sadoughi university of medical science, Yazd, Iran

Received: 25 June 2011
Accepted: 27 October 2011

Abstract

Introduction

Connection of socioeconomic status measures (such as income and education and parental addiction) to childhood leukemia are likely to vary with place and time. The aim of this study was to assess the relation between socioeconomic status and childhood leukemia.

Materials and Methods

a case- control study conducted on 86 case of acute lymphoblastic leukemia age 0-14 years in Shahid Sadoughi Hospital in Yazd and matched on age and sex to 188 healthy controls. Data was collected by interview using a questionnaire.

Data analyzed by chi-square test. Odds ratio (Ors) and 95% confidence intervals were used to measure the risk of childhood A.L.L associated with parental smoking, alcohol drinking & addiction.

Results

There was a significant difference in parental education level (P-value=0, P-value=0.001), income status (P-value =0.001), father's job (0.002) between two groups. The risk of childhood A.L.L was associated with paternal smoking (P-value =0.001, OR=2.6, CI 95%, 1.5-4.5), alcohol drinking (P-value=0.003, OR=3.33, CI 95%, 2.7-3.9), addiction (P-value =0, OR=42.7, CI95%, 5.56-328.34).

Conclusion

The results suggest that socioeconomic factors and paternal smoking and alcohol drinking are related to risk of childhood leukemia. It should be considered for planning support.

Keywords

Precursor Cell Lymphoblastic Leukemia-Lymphoma; social class; Smoking; Alcohol Drinking; Behavior, Addictive

Corresponding Author

Khadijeh Dehghani, nursing & midwifery college, Shahid Sadoughi university of medical science, Yazd, Iran, PO BOX: 89161-85366, Tel : (0351, 8241751) Fax :(0351-8249705)

Introduction

Leukemia is one of the most common potentially fatal illnesses in children (1). It accounts for approximately one third of all malignancies in this age group. Although the etiology of childhood leukemia remains undefined; a causal association with many risk factors has also been described (2).

Over the last years, the interest in assessing social inequalities and health has increased. Socioeconomic characteristics have been associated with morbidity and mortality discrepancies in many developed countries (3-5).

Relationship between social inequalities and cancer has been well studied for adults (6), but less extensively for childhood cancer and childhood leukemia seems to be unique in this aspect (7). A comprehensive review on the association between socioeconomic status (SES) and childhood leukemia was recently published and the authors have pointed out that this association is likely to vary according to time, place and study design there are more positive association in older studies and negative association in newer ones (8). Studies of the relation between parental smoking, alcohol drinking, addiction and childhood leukemia have produced inconsistent results. Cigarette smoke contains many well-established carcinogens, and both active and passive smoking have been implicated in the development of several cancers during adulthood (9). The role of parental smoking in childhood leukemia is less certain, although biologically plausible. Cigarette smoke has been linked to an increased frequency of chromosomal abnormalities (10), oxidative damage (11), and aneuploidy of sperm (12). To date several epidemiologic studies have demonstrated arguments against a strong association or even any association between maternal smoking and childhood leukemia, but others have observed a positive relationship. Reports on the effect of paternal smoking on subsequent risk of leukemia in offspring are inconclusive. Alcohol drinking is a behavior that often accompanies cigarette smoking, and has been linked to fetal growth retardation and miscarriage during pregnancy. However, the relationship between parental alcohol consumption and childhood leukemia also remain unclear. Early studies generally found no effect of maternal alcohol consumption on development of leukemia in offspring while more recent reports have observed an increased risk for both the acute lymphocytic and myeloid leukemia subtypes (9).

The authors concluded that more studies with several types of SES measures are necessary to evaluate the strength of this association. The aim of this study was to evaluate the association between socioeconomic status and childhood acute lymphoblastic leukemia.

Materials and Methods

A case-control study was conducted on 86 patients with acute lymphoblastic leukemia ages 0-14 years in Shahid Sadoughi hospital, Yazd (2008-2009). The subjects were matched on age and sex to 188 healthy controls. Controls were selected using randomized sampling among schools. The schools of control group were selected from case group. The control group included classmates and playmates of case group who were randomly selected and were the same in age and sex.

Data were collected by on interview with parents using a self-designed questionnaire, including demographic data (sex, age) and socioeconomic criteria (parental education, parental Jobs, family income and parental addiction). The levels of income were defined as: low, intermediate and high on the basis of individual record by parents. Questions were also asked about maternal and paternal smoking history, and alcohol consumption, and addiction history (prior, during pregnancy and current).

Statistical Analysis

Data were analyzed using SPSS version 15. Chi-square test was used to verify the association between acute lymphoblastic leukemia and each criteria of socioeconomic status. Odds ratio

(Odds) and 95% confidence intervals were used to measure the risk of childhood A.L.L associated with parental smoking, alcohol drinking & addiction.

Results:

62.8% (n=54) of cases and 60.1% (n=43) of controls were male. Mean age in case and control group was 7.13 ± 3.39 and 6.85 ± 3.58 . There was a significant difference in parental education level, income status, Fathers job between two groups. Of 86 patients with All, 15.1% (n=13) had illiterate fathers and 31.2% (n=26) had high school degree (diploma) or above (Academic education), While of 188 children in controls, 1.6% (n=3) had illiterate fathers and 53.8% (n=101) had high school degree or above. The difference in father's education level between two groups was significant by chi-square test (P-value <0.001) (table 1).

Illiteracy rate in patients mothers was 12.8% (n=11) in case and 3.2% (n=6) in control group 36% (n=31) of mothers in cases group had diploma or above, while it was 53.2% (n=100) in control group, which was significantly different (P-value =0.001) (table 1).

There was significant difference between fathers job in two groups (P-value =0.002) and 48.8% (n=42) of patients fathers were workers and farmers (table 2). Most of mothers in two groups were house wife. So there was no significant difference between two groups on mother's job.

Family income state in two groups was significantly different by chi-square test (P-value =0.001) (table 3). 32.6% (n=28) of cases were on low incomes while only 15.4% (n=29) of control group had such state. 7% (n=6) of cases and 13.8% (n=26) of control were placed in high income state.

Among case fathers, 43% reported having smoked, compared with 22.3% of control fathers. There was a significantly association with paternal smoking and A.L.L in children. (P-value =0.001, OR=2.625, CI 95%, 1.518-4.539) (Table 4).

Among case fathers, 5.8% reported alcohol drinking, compared with 0% of control fathers. There was a significantly association with paternal alcohol drinking and childhood all. (P-value =0.003, OR=3.33, CI=95%, 2.77-3.98) (Table 4).

Among case fathers, 18.6% reported addiction history (past and current), compared with 8% of control fathers. There was a statistically significant association between childhood A.L.L and paternal addiction (P-value <0.001 OR=42.7, CI=95%, 5.5-328.34)) (Table4).

None of mothers in two group reported alcohol drinking, addiction history and only one person in two group reported smoking. So, there was no significant association between childhood A.L.L and maternal risk factors.

Table 1: frequency distribution of mother's and father's education level

Groups		Case		Control		total	
Education level		n	%	n	%	n	%
Uneducated	mother's	11	12.8	6	3.2	17	6.2
	father's	13	15.1	3	1.6	16	5.8
Primary school	mother's	23	26.7	44	23.4	67	24.5
	father's	18	20.9	37	19.7	55	20.1
Secondary school	mother's	21	24.4	38	20.2	59	21.5
	father's	29	33.7	47	25	76	27.7
High school diploma	mother's	24	27.9	63	33.5	87	31.8
	father's	19	22.1	58	30.9	77	28.1
+above	mother's	7	8.1	37	19.7	44	16.1
	father's	7	8.1	43	22.9	50	18.2
Total	mother's	86	100	188	100	274	100
	father's	86	100	188	100	274	274
P-value	mother's	0.001					
	father's	0.000					

Table 2: frequency distribution of father's job

Groups		Case		Control		Total	
Job		n	%	n	%	n	%
Plain worker and farmer		42	48.8	53	28.2	95	34.7
State employee		12	14	53	28.2	65	23.7
Self employed		31	36	73	38.8	104	38
Non-employed		1	1.2	9	4.8	10	3.6
Total		86	100	188	100	274	100

P-value =0.002

Table 3: frequency distribution of family income state

Groups		Case		Control		Total	
Income state		N	%	n	%	n	%
Low		28	32.6	29	15.4	57	20.8
Intermediate		52	60.4	133	70.7	185	67.5
High		6	7	26	13.8	32	11.7
Total		86	100	188	100	274	100

P-value =0.001

Table4: Frequency distribution of paternal smoking, alcohol consumption and addiction

Parameter	Group	Case		Control		Sum		P-value	OR
		Yes	No	Yes	No	Yes	No		
Smoking	n	37	49	42	146	195	79	0.001	2.625
	%	43	57	22.3	77.7	71.2	28.8		
Alcohol	n	5	81	0	188	5	269	0.003	3.33
	%	5.8	94.2	0	100	1.8	98.2		
Addiction	n	16	70	1	187	17	257	0.000	42.7
	%	18.6	81.4	5	99.5	6.2	93.8		

Discussion

In this study there was a negative association between childhood acute lymphoblastic leukemia and family income, parental education level and father's job .Several studies have shown an association between leukemia and high SES (13-18). Over all, studies utilizing area based socioeconomic measures have demonstrated an increased risk of A.L.L. among people with high SES (7, 1, and 19). On the other hand, the results of the studies of SES and childhood leukemia using individual level assessment are controversial (15-17, 20-24). High levels of family income and parental education, reassured individually have been consistently associated with a lower risk of childhood leukemia, while association of parental occupational class with childhood leukemia demonstrates a contrary association; i.e. high rates are correlated with high SES (49-51), Including findings from 2 cohort studies (25-26).A recent case-control study conducted in united kingdom did not show any difference in childhood A.L.L risk according to deprivation levels, whether using area-or. Individual based measure of SES at the time of birth or diagnosis (27).Recently, Pool, et al. pointed out the difficulties in making quantitative comparisons between studies, since many different types of SES measure were utilized and their distinct social implications can vary by place and time (8). In fact, the adequacy of the measure is also related to the study aims and availability of information in each particular country (28). In this study, high percent of fathers in case group were workers or farmers, it can indicates social class of job and probably shows that fathers contact with risk agent at work can increase risk of childhood A.L.L which has been reported in several studies (29) We categorized family income state as low, intermediate, and high based on the parents view, and amount of family income and other factors such as family size, living place, and social class of parental job have not been recorded. It is recommended to conduct cohort studies with more subjects, and more accurate SES measures.

In agreement with our results several studies have found a positive relationship between childhood all and paternal smoking. (30-34) and some others is controversial (37, 38, 9, 35). Certainly, more studies are needed to better understand the effect of parental smoking on childhood A.L.L risk. (30). It is recognized that childhood A.L.L and AML (acute meylcytic leukemia) are two distinct diseases with different histological presentations, age distributions, and prognoses. (39)

Our results were consistent with a few studies that reported relationship between paternal alcohol drinking and childhood leukemia (30).

And several studies reported controversial (9, 31).Alcohol metabolism is known to produce reactive oxygen species (ROs) that could contribute to carcinogenesis (38).

According to the results of this study, paternal addiction had a positive association with acute childhood leukemia. Yet, any studies have done about this relationship. In our study, mothers didn't have history of (%1), alcohol addiction and cigarette smoking was equal (%1) in two groups. Many studies have found a positive relationship between childhood leukemia and maternal alcohol consumption (9, 31) although the evidence is inconclusive (40). It is unclear whether maternal or paternal cigarette smoking before or during pregnancy is a risk factor for developing childhood leukemia (30, 35).

Conclusion

According to this study findings, child-hood leukemia showed to be related with families low SES. Thus, it is highly recommended to consider supportive plans for these patients to improve treatment outcomes.

Acknowledgement

The authors gratefully acknowledge the nursing staff in pediatric ward of Shahid Sadoughi in hospital in Yazd, for their help, and MS. Maryam Nikfard for editing the full text article English draft.

Conflict of Interest

This study was supported by Shahid Sadoughi University of Medical Sciences Yazd, Iran.

References

1. Borugian marilyn J, Spinellin Jjohn J, Mezel Gabor, Wilkins Russell, Abanto Zenadia, Mcbride mary L. childhood leukemia and socioeconomic status in Canada; epidemiology. 2005, 16(4), 526-31.
2. Karina Braga ribeiro, Patricia A. Buffer, Catherine netaye. Socioeconomic status and childhood acute lymphocytic leukemia incidence in Sao Paulo, Brazil. International Journal of Cancer, 2008; 123(8), 1907-1912.
3. Sorlie PD, Backlund E, Keller JB. Us mortality by economic, demographic, and social characteristics: the national long it udinal mortality study. Am J Public Health 1995; 85: 949-56.
4. Adler NE, Ostrove JM. Socioeconomic status and health: what we know and what we don't Ann N Y Acad Sci 1999; 896: 3-15.
5. Singh Gk. Area deprivation and widening inequalities in us mortality, 1969-1998. Am J Public Health 2003; 93: 1137-43.
6. Kogevians M, pearce N. Susser M, Boffetta P, eds. Social inequalities and cancer. Lyon: international agency for research on cancer, 1997.
7. Greenberg Rs, Shuster JL. Epidemioloty of cancer in children. Epidemiol Rev 1985; 7: 22-48.
8. Pool C, Green Land S, Luetters C, Kelsey JL, Mezei G. socioeconomic status and childhood leukemia: a review. Int J. Epidemiol 2006; 35: 370-840.
9. Macarthur C. Amy, Mcbrid L. Mary, Spinelli Y. John, Tamaro Sharon, Gallagher P. Richard, theriault gilles, cancer causes control. 2008; 19: 238-295.
10. Pluth Jm, Ramsey My, Tucker JD. Role of maternal exposures and newborn genotypes on newborn chromosome aberration frequencies. Mutat Res. 2000; 465: 101-111.
11. Fraga CG, Motchnik PA, Wyrobek AJ, Rempel PM, Ames BN. Smoking and low antioxidant levels increase oxidative damage to sperm DNA. Mutal Res. 1996; 351: 199-203.
12. Shi A, Ko E, Barclay L, Hoang T, pademaker A. Martin R. cigarette smoking and aneuprordy in human sperm. Mol reprod Dev. 2001, 59: 417-421.
13. Alexander FE, Carwright RA, Mc Kinney PA, Cartwright RA. Community lifestyle characteristics and risk of acute lymphoblastic leukemia in children. Lance 1990; 336: 1461-5.
14. Alexander FE, Cartwright RA, Mckinney, PA, Ricketts Ty. Leukemia incidence, social class and estuaries: an ecological analysis. J public health Med 1990; 12: 109-17.
15. Kuehni CE, Zwalen M. commentary: Numerous, heterogeneous, and often poor- the stuies on childhood leukemia and socioeconomic status. Int J Epidemiol 2006; 35: 385-5.
16. Petridou E, Kassimos D, Kalmanti M, Kasmidis H, Haidas S, Flytzani V, et al. Age of exposure to infections and risk of childhood leukaemia. BMJ 1993; 304-774.

17. Petridou E, Trichopoulos D, Kalapthaki V, Pourtsidis A, Kogerinas M, Kalmanti M , et al. the risk profile of childhood leukemia in crece: a nationwide case-control study Br J cancer 1997; 76: 1241-7.
18. Li Cy, Lin RS, Lin CH. Urbanization and childhood leukemia in Taiwan. Int J Epidemiol 1998; 27: 587-91.
19. Parslow RC, Law GR, Feltbower R, Kinsey SE, MCKinney PA. population mixing, childhood leukemia CNS tumors and other childhood cancers in. York shire. Eur J cancer 2002; 38: 2033-40.
20. Rejnolds P, Von Behren J, Elkin Ep. Birth charac teristics and leukemia in young children. Am J Epiemiol 2002; 155: 603-13.
21. Brondum J, Shu Xo, Steinbuch M. Severson RK, Potter Jd, Robinson LL. Parental cigarette smoking and the risk of acute leukemia in children. Cancer 1999; 85: 1380-8.
22. Dockerty JD, Skegg DC, Elwood JM, Herbison Gp, Becroft DM, Lewis ME. Infections, vaccinations, and the risk of childhood leukemia. Br J cancer 1999; 80: 1483-9.
23. Raaschou-Nielsen O, Obel J, Dalton S, T Jonne land A, Hansen J. socioeconomic status and risk of childhood leukemia in Denmark. Scand J public Health 2004; 32: 279-86.
24. Wong DL, Dockery JD. Birth characteristic S and the risk of childhood leukemia and lymphomas in New Zealand: a case-control study. BMC Blood Disord 2006; 6: 5-11.
25. Dickinson Ho. Parker L. Quantifying the effect of population mixing on childhood leukemia risk: the seascale cluster. Br cancer 1999; 81: 144-51.
26. Murray L, Mc Carron P, Bailie K, Middleton R, Davey Smith G, Dempsey S. and et al. association of early life factors and acute lymphoblastic leukemia in childhood: historical cohort study. Br J cancer 2002; 86: 356-61.
27. Smith A, Roman E, Simpson J, Ansell P, fear NT, Eden T. childhood leukaemia and socioeconomic stutus: fact or artifact? A report from the United Kingdom childhood cancer study (UKCCS). Int J Epidemiol 2006; 35: 1504-13.
28. Berkman LF, Macintyre S. the measurement of social class in health studies: old measures and new formulations. In: Kogevinas M, pearce N. Susser M, Boffetta P, editors. Social inequalities and cancer. Lyon: international agency for research of cancer, 1997, 51-64.
29. Belso M, Kingsley B, Holmes A. Risk factors for acute leukemia in children a review. Environ health perspect. 2007; 115(1): 138-45.
30. Shu Xo, Ross YA, Pendergrass Tw, Reman GH, Lampkin B, Rabison LL. Parental alcohol consumption, cigarette smoking, and risk of infant leukemia; a childrens cancer group study. J Natl cancer Inst, 1996, 88: 2431.
31. Sorahan T, Lancashire R, prior P, Peck I, Stewart A. childhood cancer and parental use of alcohol and tobacco. Ann Epidemiol, 1995; 5: 354-459.
32. John EM, Savitz DA, Sandler DP. Parental exposure to parents smoking and childhood cancer. Am J epidemiol; 1991, 133: 123-132.
33. Ji BT, Shu Xo, Linet MS and etal. Parental cigarette smoking and risk of childhood cancer among offspring of nonsmoking mothers. Y Natl cancer Inst, 1997; 889: 238-244.
34. Chang Yeffrey S, Selvin Steve, Metayer Catherine, crouse Vonda, Golembesky Amanda and buffler A. patricia. Parental smoking and the risk of childhood leukemia, American Journal of epidemiology, 2006, 163 (12): 1091-1100.
35. Brondum Y, Shu Xo, Stein buch M. Parental cigarette smoking and the risk of acute leukemia in children cancer, 1999; 85: 1380-88.
36. Pang D, mC Nany R, Birch YM. Parental smoking and childhood cancer, Br Y cancer, 2003; 88: 373-81.
37. Annie J sascos Harri vaninio. Parental smoking to childhood cancer: a possible link and the need for action, human & experimental toxicology; 1999, 18(4): 192-201
38. Magnan C, pastor G, Luzzato L. Parental occupation and other environmental factors in the etiology of leukemais and non-hodgkins lymphomas in childhood: a case-control study. Tumori, 1990; 76: 413-19.
39. Infante- Rivard C, krajnovic M, Labuda D, et al. Parental smoking, CYP1A1 genetic polymorphism and childhood leukemia. Epidemiology. 2002; 13: 277-81.
40. Ries LAG, Smith MA, Gurney YG. Cancer incidence and survival among children and adolescents. United States SEER program. National cancer institute; 1999, 4049