Effect of 5-fluoro-2'-deoxycytidine (FdCyd) on p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b Genes Expression, Apoptosis Induction, and Cell Growth Inhibition in Pancreatic Cancer AsPC-1 and Hepatocellular Carcinoma LCL-PI 11 Cell Lines

Masumeh Sanaei MSc1, Fraidoon Kavoosi PhD1,*

1. Research center for non-communicable diseases, Jahrom University of medical sciences, Jahrom, Iran *Corresponding author: Dr Fraidoon Kavoosi, Jahrom University of Medical Sciences, Jahrom, Fars province, Iran. Email: kavoosifraidoon@gmail.com. ORCID ID: 0000-0001-7761-7912

Received: 07 June 2020 Accepted: 25 October 2020

Abstract

Background: Aberrant DNA methylation of the promoter region is one of the most epigenetic changes in numerous cancers. DNA methyltransferase inhibitors (DNMTIs) can revert DNA hypermethylation in tumor suppressor genes (TSGs). The present study was designed to investigate the effect of 5-fluoro-2'-deoxycytidine (FdCyd) on p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b genes expression, apoptosis induction, cell growth inhibition in pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines.

Materials and Methods: The cells were treated with FdCyd at different periods. Then, the MTT assay, cell apoptosis assay, and qRT-PCR were done to determine cell viability, cell apoptosis, and the relative gene expression level respectively.

Results: The FdCyd decreased DNA methyltransferase 1, 3a, and 3b and increased p16INK4a, p14ARF, and p15INK4b genes expression significantly (P<0.001). Besides, LCL-PI 11 cell was more sensitive to FdCyd in comparison to AsPC-1 cell. FdCyd induced significant cell growth inhibition with a dose- and time-dependent manner (P<0.001). The IC50 value of FdCyd was obtained with approximately 1 μ M. Further, FdCyd induced cell apoptosis significantly as a time-dependent manner. The number of apoptotic cells was significantly increased in all groups. The percentage of apoptotic cells after 24 and 48 h were 13.86 and 29.6 % in AsPC-1 and 21.04 and 41.52 % in LCL-PI 11 cell line respectively (P<0.001).

Conclusion: The FdCyd can reactivate the p16INK4a, p14ARF, and p15INK4b through inhibition of DNA methyltransferase 1, 3a, and 3b gene expression.

Keywords: 5-fluoro-2'-deoxycytidine, Genes, Neoplasm, Onco-Suppressor

Introduction

The mammalian cell cycle is controlled by molecular mechanisms and several pathways ensuring correct progression and cell division. This division is divided into two stages, mitosis and interphase leads to DNA replication and segregation of replicated chromosomes into two new cells. The interphase comprises G1, S, and G2 phases (1). To date, many studies have reported several genetic and epigenetic alterations of cell-cycle regulators in various cancers. DNA hypermethylation of tumor suppressor genes (TSGs) such as the INK4 family has been indicated in humans, the numerous cancers. In

INK4b/ARF/INK4a gene locus located on chromosome 9p21 and encodes potent TSGs including p15ink4b, p14ARF, and p16ink4a. The p15ink4b and p16ink4a bind to CDK4 and 6 and act as cell cycle inhibitors in the G1 phase cell cycle (2). Global cytosine methylation patterns of the promoter region in mammalian cells are established by a complex interplay of at least three independently encoded DNA methyltransferases (DNMTs) consists of DNMT1, DNMT3A, and DNMT3B (3). The levels of these enzymes are often increased in numerous cancer cell lines, which may account for the DNA methylation of TSGs in numerous cancers

(4). Hypermethylation of CpG islands of TSGs may be involved in carcinogenesis in gastric cancer, breast cancer, pancreatic cancer, head and neck cancer, hepatic cancer, lung cancer, colorectal cancer, bladder cancer, and renal cancer (5). Several compounds act as the DNA methyltransferase inhibitor (DNMTI) 5-Aza-2'-deoxycytidine (5comprising 5-azacytidine AZA-CdR, Decitabine), (azacitidine), 1-beta-D-arabinofuranosyl-5dihydro-5azacytosine (fazarabine), azacytidine (DHAC), 5-fluoro-2'deoxycytidine (FdCyd) (6) and zebularine (7). Previously, we reported the effect of 5-AZA-CdR on hepatocellular carcinoma (HCC) (8, 9) and colon cancer (10). The present study aimed to investigate the effect of 5-fluoro-2'-deoxycytidine p16INK4a, p14ARF. (FdCyd) on p15INK4b, and DNA methyltransferase 1, 3a, and 3b genes expression, apoptosis, cell growth inhibition induction in pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines.

Materials and Methods Materials

Human pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines were purchased from the National Cell Bank of Iran-Pasteur Institute. FdCyd and Dulbecco's modified Eagle's medium (DMEM) were obtained from Sigma (St. Louis, MO, USA). The compound was dissolved in dimethyl sulfoxide (DMSO) to make a work stock solution. Further concentrations of the agent were obtained by diluting the provided solution. Other compounds including materials and various kits were purchased as provided for previous works (11, 12). The cells were maintained in DMEM supplemented with fetal bovine serum 10% and antibiotics in a humidified atmosphere of 5% CO2 in air at 37°C. This work was approved in the Ethics Committee of Jahrom University of Medical science with a code number of IR.JUMS.REC. 1399.009.

Cell culture and cell viability

Human pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines were cultured in DMEM supplemented with 10% FBS and antibiotics at 37°C in 5% CO2 overnight and then seeded into 96-well plates (3×10^5) cells per well). After 24 h, the medium was replaced with an experimental medium containing various concentrations of 5fluoro-2'-deoxycytidine (0, 0.5, 1, 2.5, 5, and 10 µM). The control groups received DMSO only, at a concentration of 0.05 %. After a period of 24 and 48h, the cells were investigated by MTT assay according to Standard protocols to determine cell viability. Therefore, MTT solution was added to each well for 4 h at 37°C and then the MTT solution was changed by DMSO and shaken for 10 min to dissolve all of the crystals. Finally, the optical density was detected by a microplate reader at a wavelength of 570 nM. Each experiment was repeated three times (triplicates).

Cell apoptosis assay

To determine AsPC-1 and LCL-PI 11 apoptotic cells, the cells were cultured (3 \times 10⁵ cells/well) and incubated overnight and then treated with FdCyd (1 µM) for different time periods (24 and 48 h). Subsequently, the treated and untreated cells were harvested by trypsinization, washed with cold PBS, and resuspended in Binding buffer (1x). Finally, Annexin-V-(FITC) and PI were used to determine the apoptotic cells FACScan by flow cytometry (Becton Dickinson, Heidelberg, Germany).

Real-time Quantitative Reverse Transcription Polymerase

Chain Reaction (qRT-PCR)

To determine the relative expression level of p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b genes qRT-PCR was done. The AsPC-1 and LCL-PI 11 cells were treated with FdCyd (1 μ M) for 24 and 48 h and then qRT-PCR was done as our previous works (13). The primer sequences are indicated in Table I.

2

DOI: 10.18502/ijpho.v11i1.5000

Statistical analysis

The database was set up with the SPSS 16.0 software package (SPSS Inc., Chicago, Illinois, USA) for analysis. The data were acquired from three separate tests and are indicated as means \pm standard deviations. The comparisons between groups were achieved with ANOVA (one-way ANOVA) and Turkey test. A significant difference was considered as P < 0.05.

Results

Result of cell viability by the MTT assay The cell viability of the AsPC-1 and LCL-PI 11 cells treated with FdCyd (0, 0.5, 1, 2.5, 5, and 10 μ M) was investigated by MTT assay. This technique showed that the activity of cellular enzymes reduced the tetrazolium salt MTT and produced a dark-blue formazan crystal. The crystal was dissolvable in DMSO by which the number of viable cells can be investigated. As indicated in Figure. 1, FdCyd induced significant cell growth inhibition with a dose-dependent manner (P< 0.001). The IC50 value of FdCyd was obtained with approximately 1 μ M.

Result of cell apoptosis assay

To determine apoptosis, the pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cells were treated with FdCyd (1 μ M) for 24 and 48 h and then stained using annexin-V-(FITC) and PI to determine apoptotic cell in early and late apoptosis stage. As indicated in Figures 2 and 3, FdCyd induced cell apoptosis significantly as a time-dependent manner in all experimental groups (P<0.001).

Result of determination of genes expression

The effect of FdCyd (1 μ M) on p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b genes expression was investigate by quantitative real-time RT-PCR analysis. The result indicated that treatment with FdCyd (1 μ M) up-regulated p16INK4a, p14ARF, p15INK4b, and down-regulated DNA methyltransferase 1, 3a, and 3b genes expression significantly, Figures 4 and 5 and Table II. FdCyd had a significant time-dependent manner (P<0.001).

Table I. The primer sequences of p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3	3a,
and 3b genes.	

	e	
Primer	Primer sequences (5' to 3')	Reference
DNMT1		13
Forward	GAGGAAGCTGCTAAGGACTAGTTC	
Reverse	ACTCCACAATTTGATCACTAAATC	
DNMT 3a		13
Forward	GGAGGCTGAGAAGAAAGCCAAGGT	
Reverse	TTTGCCGTCTCCGAACCACATGAC	
DNMT 3b		13
Forward	TACACAGACGTGTCCAACATGGGC	
Reverse	GGATGCCTTCAG GAATCACACCTC	
p14 ^{ARF}		14
Forward	GTGGGTTTTAGTTTGTAGTT	
Reverse	AAACCTTTCCTACCTAATCT	
p15INK4b		15
Forward	AAGCTGAGCCCAGGT CTCCTA	
Reverse	CCACCGTTGGCCGTAAACT	
p16INK4a		16
Forward	CCCGCTTTCGTAGTTTTCAT	
Reverse	TTATTTGAGCTTTGGTTCTG	
GAPDH		12
Forward	AAC GTG TCA GTO GTG GAC CTG	
Reverse	GGG TGT CGC TGT FGA AGT	

Iran J Ped Hematol Oncol. 2021, Vol 11, No 1, 01-10

Cell line	Gene	Drug		Duration (h)	Expression	P-value
- DC 4	1 (D) 11/4	F1G 1	(μΜ)	24		D :0.001
AsPC-1	p16INK4a	FdCyd	1	24	2.3	P<0.001
AsPC-1	p16INK4a	FdCyd	1	48	2.7	P<0.001
AsPC-1	p14ARF	FdCyd	1	24	1.8	P<0.001
AsPC-1	p14ARF	FdCyd	1	48	2.3	P<0.001
AsPC-1	p15INK4b	FdCyd	1	24	2.1	P<0.001
AsPC-1	p15INK4b	FdCyd	1	48	2.5	P<0.001
AsPC-1	DNMT1	FdCyd	1	24	0.54	P<0.001
AsPC-1	DNMT1	FdCyd	1	48	0.47	P<0.001
AsPC-1	DNMT3a	FdCyd	1	24	0.55	P<0.001
AsPC-1	DNMT3a	FdCyd	1	48	0.46	P<0.001
AsPC-1	DNMT3b	FdCyd	1	24	0.48	P<0.001
AsPC-1	DNMT3b	FdCyd	1	48	0.42	P<0.001
LCL-PI 11	p16INK4a	FdCyd	1	24	2.6	P<0.001
LCL-PI 11	p16INK4a	FdCyd	1	48	2.7	P<0.001
LCL-PI 11	p14ARF	FdCyd	1	24	2.1	P<0.001
LCL-PI 11	p14ARF	FdCyd	1	48	2.3	P<0.001
LCL-PI 11	p15INK4b	FdCyd	1	24	2.5	P<0.001
LCL-PI 11	p15INK4b	FdCyd	1	48	2.8	P<0.001
LCL-PI 11	DNMT1	FdCyd	1	24	0.46	P<0.001
LCL-PI 11	DNMT1	FdCyd	1	48	0.34	P<0.001
LCL-PI 11	DNMT3a	FdCyd	1	24	0.42	P<0.001
LCL-PI 11	DNMT3a	FdCyd	1	48	0.39	P<0.001
LCL-PI 11	DNMT3b	FdCyd	1	24	0.39	P<0.001
LCL-PI 11	DNMT3b	FdCyd	1	48	0.32	P<0.001

Table II. The relative expression level of p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase
1, 3a, and 3b genes.



B: LCL-PI cell

Figure 1. In vitro effects of FdCyd (0, 0.5, 1, 2.5, 5, and 10 μ M) on AsPC-1 (A) and LCL-PI 11 (B) cell viability determined by MTT Assay at 24 and 48 h. The FdCyd inhibited the growth of both cell lines significantly as a dose-dependent manner.



Figure 2. The apoptotic effect of FdCyd (1 μ M) on AsPC-1 cell versus control groups at different periods (24 and 48h). The FdCyd induced significant apoptosis as a time-dependent manner. The results were obtained from three separate experiments and were presented as mean \pm standard error of the mean.



Figure 3. The apoptotic effect of FdCyd (1 μ M) on LCL-PI 11 cell versus control groups at different periods (24 and 48h). The FdCyd induced significant apoptosis as a time-dependent manner. The results were obtained from three separate experiments and were presented as mean \pm standard error of the mean.



Figure 4. The relative expression level of p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b in the pancreatic cancer AsPC-1 cell line treated with FdCyd (1 µM) versus untreated control groups at different periods (24 and 48h). 0: Control, 1: 24 h, 2: 48 h.



Figure 5. The relative expression level of p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b in the hepatocellular carcinoma LCL-PI 11 cell line treated with FdCyd (1 µM) versus untreated control groups at different periods (24 and 48h). 0: Control, 1: 24 h, 2: 48 h.

6

Discussion

of the Aberrant DNA methylation promoter region is one of the most epigenetic changes in numerous cancers. The activity of DNMTs leads to DNA hypermethylation and cancer induction (17). Meanwhile, the methylation of CpG islands affects the transcription and expression of specific genes such as p16INK4a, p14ARF, p15INK4b leads to cancer induction. However, DNMTIs can revert DNA hypermethylation in tumor suppressor genes (TSGs) and suppress cancer-specific cellular phenotypes (18). In the current study, we indicated that down-regulated DNA FdCyd methyltransferase 1, 3a, and 3b genes up-regulated expression. p16INK4a. p14ARF, and p15INK4b, inhibit cell growth and induced apoptosis in pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines. Similarly, our previous studies indicated that DNA demethylating agent 5aza-CdR (3 µM) can induce apoptosis by p16INK4a, p14ARF, p15INK4b genes reactivation in HCC, PLC/PRF5, and pancreatic cancer, MIA Paca-2, cell lines (9). In line with our report, it has been demonsterated that DNA demethylating agent 5-Aza-CdR can induce p14ARF mRNA expression in esophagus cancer cell lines (TE-1 and Eca-9706) (19), p16INK4a in pancreatic cancer cell lines and SW1990 AsPC-1 (20).and P15/INK4b and P16/INK4a in acute myeloid leukemia (AML) KBM3/Bu2506, OCI-AML3, and HL60 cell lines (21). We did further investigation and found that FdCyd played its role through the downregulation of DNA methyltransferase 1, 3a, and 3b gene expression. Inconsistent with our results, in vitro study has shown that zebularine [1-(beta-D-ribofuranosvl)-1, 2-dihvdropyrimidin-2-one] reactivates a silenced p16INK4a gene and demethylates its promoter region in T24 bladder cancer cell line (22). It has been shown that DNA methyltransferase inhibitor 5-aza-2'-

deoxycytidine (decitabine) caused the DNA demethylation of the all CpG sites in p15INK4b islands the CpG in hematopoietic cells (23). Furthermore, other researchers have reported that FdCyd induces p21 gene reactivation which correlates with the arrest of AML cells in the G_1 cell cycle phase (24). It has also shown been that the p21^{WAF1/CIP1} expression in metastatic prostate cancer PC3, LNCaP, and DU145 cell lines is enhanced as a result of treatment with 5-Aza-CdR (25). Besides, 5-Aza-dC can increase the up-regulation of p21^{WAF1}, p27^{KIP1}, and p57^{KIP2} genes expression in pancreatic cell lines AsPC-1, PANC-1, BxPC-3, HPAC, Capan-2, and (26). Our previous work HPAF-II indicated that 5-AZA-CdR can downregulate DNMT1, DNMT3a, DNMT3b in HCC LCL-PI 11 cell line [27]. Furthermore, we indicated that 5-Aza-CdR can induce apoptosis through the reactivation of p15INK4, p16INK4, p18INK4, and p19INK4 genes expression in HCC PLC/PRF/5 cell line [28]. It has reported that 5-Aza-CdR been significantly increases the expression of P16 leads to apoptotic induction in pancreatic cancer [29]. In T24 human bladder carcinoma, it has been shown that 5-Aza-CR and zebularine induce apoptosis by up-regulation of p16 [30]. Finally, DNA demethylating agents can reactivate cyclin-dependent kinase inhibitors by inhibition of DNA methyltransferase inhibitors activity. It should be noted that our report is the first report of the evaluation of the effect of FdCyd on p16INK4a, p14ARF, p15INK4b, and DNA methyltransferase 1, 3a, and 3b genes expression, apoptosis, and cell growth inhibition induction in pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines. Therefore, it is the novelty of the current work. We did not assess the effect of the compound at 72 h because of budget limitation. This evaluation is recommended.

Conclusion

In summary, our findings indicated that FdCyd can reactivate the p16INK4a, p14ARF, p15INK4b and through inhibition of DNA methyltransferase 1, 3a, and 3b genes expression in pancreatic cancer AsPC-1 and hepatocellular carcinoma LCL-PI 11 cell lines resulting in cell growth inhibition and apoptosis induction. Further, the LCL-PI 11 cells were more sensitive to FdCyd in comparison to the pancreatic cancer AsPC-1 cells.

Acknowledgement

This article was supported by the adjutancy of research of Jahrom University of Medical Sciences, Iran.

Conflict of interest

The authors declare no conflict of interest.

References

1. Vermeulen K, Van Bockstaele DR, Berneman ZN. The cell cycle: a review of regulation, deregulation and therapeutic targets in cancer. Cell Prolif 2003; 36(3):131-149.

2. Ko A, Han SY, Song J. Regulatory network of ARF in cancer development. Mol Cells 2018; 41(5): 381-389

3. Robertson KD. DNA methylation, methyltransferases, and cancer. Oncogene 2001; 20(24):3139-3155.

4. Subramaniam D, Thombre R, Dhar A, Anant S. DNA methyltransferases: a novel target for prevention and therapy. Front Oncol 2014; 4:80-86.

5. Daniel FI, Cherubini K, Yurgel LS, de Figueiredo MAZ, Salum FG. The role of epigenetic transcription repression and DNA methyltransferases in cancer. Cancer 2011; 117(4):677-687.

6. Goffin J, Eisenhauer E. DNA methyltransferase inhibitors—state of the art. Ann Oncol 2002;13(11):1699-1716.

7. Lyko F, Brown R. DNA methyltransferase inhibitors and the development of epigenetic cancer therapies. J Natl Cancer Inst 2005; 97(20):1498-1506.

8. Sanaei M, Kavoosi F. Effects of 5-aza-2'-deoxycytidine and Valproic Acid on Epigenetic-modifying DNMT1 Gene Expression, Apoptosis Induction and Cell Viability in Hepatocellular Carcinoma WCH-17 cell line. Iran J Ped Hematol Oncol 2019; 9 (2): 83-90.

9. Sanaei M, Kavoosi F, Mohammadi M, Khanezad M. Effect of 5-aza-2'deoxycytidine on p16INK4a, p14ARF, p15INK4b Genes Expression, Cell Viability, and Apoptosis in PLC/PRF5 and MIA Paca-2 Cell Lines. Iran J Ped Hematol Oncol 2019; 9(4): 219-228.

10. Sanaei M, Kavoosi F. Effect of 5-aza-2'-deoxycytidine in comparison to valproic acid and trichostatin A on histone deacetylase 1, DNA methyltransferase 1, and CIP/KIP family (p21, p27, and p57) genes expression, cell growth inhibition, and apoptosis induction in colon cancer SW480 cell line. <u>Adv Biomed Res</u> 2019; 8: 52-58.

11. Sanaei M, Kavoosi F, Roustazadeh A, Shahsavani H. In vitro effect of the histone deacetylase inhibitor valproic acid on viability and apoptosis of the PLC/PRF5 human hepatocellular carcinoma cell line. APJCP 2018;19(9):2507-2512.

12 Sanaei M, Kavoosi F. Effect of curcumin and trichostatin a on the expression of DNA methyltransfrase 1 in hepatocellular carcinoma cell line hepa 1-6. Iran J Ped Hematol Oncol 2018; 8(4):193-201.

13. Sanaei M, Kavoosi F, Roustazadeh A, Golestan F. Effect of genistein in comparison with trichostatin a on reactivation of DNMTs genes in hepatocellular carcinoma. J Clin Transl Hepatol 2018; 6(2):141–146.

14. Sakuma K, Chong JM, Sudo M, Ushiku T, Inoue Y, Shibahara J, et al. High-density methylation of p14ARF and p16INK4A in Epstein-Barr virus– associated gastric carcinoma. Int J Cancer 2004; 112(2):273-278. 15. Li G, Ji Y, Liu C, Li J, Zhou Y. Reduced levels of p15INK4b, p16INK4a, p21cip1 and p27kip1 in pancreatic carcinoma. Mol Med Report 2012; 5(4):1106-1110.

16. Saegusa M, Machida B D, Okayasu I. Possible associations among expression of p14ARF, p16INK4a, p21WAF1/CIP1, p27KIP1, and p53 accumulation and the balance of apoptosis and cell proliferation in ovarian carcinomas. Cancer 2001; 92:1177–1189.

17. Kanai Y, Ushijima S, Kondo Y, Hirohashi Y, S. Nakanishi DNA methyltransferase expression and DNA methylation of CPG islands and peri-centromeric satellite regions in human colorectal and stomach cancers. Int J Cancer 2001; 91(2):205-212.

18. Stresemann C, Brueckner B, Musch T, Stopper H, Lyko F. Functional diversity of DNA methyltransferase inhibitors in human cancer cell lines. Cancer Res 2006; 66(5):2794-2800.

19. Ling Y, Zhang C, Shen R, Xu Y, Zhu C, Lu M, et al. p14ARF repression induced by promoter methylation associated with metastasis in esophageal squamous cell carcinoma. Dis Esophagus 2014; 27(2):182-187.

20. Han T, Zhuo M, Hu H, Jiao F, Wang L-W. Synergistic effects of the combination of 5-Aza-CdR and suberoylanilide hydroxamic acid on the anticancer property of pancreatic cancer. Oncol Rep 2018; 39(1):264-270.

21. Valdez BC, Li Y, Murray D, Ji J, Liu Y, Popat U, et al. Comparison of the cytotoxicity of cladribine and clofarabine when combined with fludarabine and busulfan in AML cells: Enhancement of cytotoxicity with epigenetic modulators. Exp Hematol 2015; 43(6):448-461. e2.

22. Cheng JC, Matsen CB, Gonzales FA, Ye W, Greer S, Marquez VE, et al. Inhibition of DNA methylation and reactivation of silenced genes by zebularine. J Natl Cancer Inst 2003; 95(5):399-409. 23. Zhao XY, Sakashita K, Kamijo T, Hidaka E, Sugane K, Kubota T, et al. Granulocyte–macrophage colonystimulating factor induces de novo methylation of the p15 CpG island in hematopoietic cells. Cytokine 2005; 31(3):203-212.

24. Schmelz K, Wagner M, Dörken B, Tamm I. 5-Aza-2'-deoxycytidine induces p21WAF expression by demethylation of p73 leading to p53-independent apoptosis in myeloid leukemia. Int J Cancer 2005; 114(5):683-695.

25. Bott S, Arya M, Kirby R, Williamson M. p21 WAF1/CIP1 gene is inactivated in metastatic prostatic cancer cell lines by promoter methylation. Prostate Cancer Prostatic Dis 2005; 8(4):321-326.

26. Kumagai T, Wakimoto N, Yin D, Gery S, Kawamata N, Takai N, et al. Histone deacetylase inhibitor, suberoylanilide hydroxamic acid (Vorinostat, SAHA) profoundly inhibits the growth of human pancreatic cancer cells. Int J Cancer 2007; 121(3):656-665.

27. Sanaei M, Kavoosi F, Esmi Z. The Effect of 5-Aza-2'-Deoxycytidine in Combination to and in Comparison with Vorinostat on DNA Methyltransferases, Histone Deacetylase 1, Glutathione S-Transferase 1 and Suppressor of Cytokine Signaling 1 Genes Expression, Cell Growth Inhibition and Apoptotic Induction in Hepatocellular LCL-PI 11 Cell Line. Int J Hematol Oncol Stem Cell Res 2020; 14(1): 45–55.

28. Sanaei M, Kavoosi F, Ghasemi A. Investigation of the effect of 5-Aza-2'deoxycytidine on p15INK4, p16INK4, p18INK4, and p19INK4 genes expression, cell growth inhibition, and apoptosis induction in hepatocellular carcinoma PLC/PRF/5 cell line. Adv Biomed Res 2020;9(1):33-39.

29. Han T, Zhuo M, Hu H, Jiao F, Wang L-W. Synergistic effects of the combination of 5-Aza-CdR and suberoylanilide hydroxamic acid on the anticancer property of pancreatic cancer. Oncol Rep2018;39(1):264-270.

30. Cheng JC, Matsen CB, Gonzales FA, Ye W, Greer S, Marquez VE, et al. Inhibition of DNA methylation and reactivation of silenced genes by zebularine. J Natl Cancer Inst 2003; 95:399–409.

[Downloaded from ijpho.ssu.ac.ir on 2025-07-31]