A Comparison of the Effect of Chlorhexidine Mouthwash and Combined Mouthwash (4 drugs) in Preventing and Managing Oral Mucositis in Neutropenic Patients Admitted to the Pediatric Oncology Department: A Randomized Clinical Trial

Roohollah Edalatkhah MD^{1, 2, 3}, Maryam Sadat Yazdanparast MD^{1, 2}, Motahareh Taghvaei MD^{*2}

1. Hematology and Oncology Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

2. Department of Pediatrics, Shahid Sadoughi Hospital, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

3. Children Growth Disorder Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

*Corresponding author: Dr. Motahareh Taghvaei, Department of Pediatrics, Shahid Sadoughi Hospital, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. Email: motahareh.taghvaei.73@gmail.com. ORCID ID: 0000-0002-3436-1424.

Received: 01 November 2024 Accepted: 15 March 2025

Abstract

Background: Given the lack of comprehensive studies comparing the effects of chlorhexidine and combined mouthwashes in preventing oral mucositis in our country, particularly in Yazd province, this study aims to evaluate the effectiveness of combined mouthwash versus chlorhexidine mouthwash in preventing oral mucositis in neutropenic patients.

Materials and Methods: This study was a double-blind randomized controlled trial conducted on children undergoing chemotherapy at Shahid Sadoughi Hospital. Patients were randomly assigned to two groups of 30: one group received combined mouthwash containing diphenhydramine, nystatin, aluminum-magnesium hydroxide, and lidocaine (group 1), while the other group received a 0.2% chlorhexidine mouthwash (group 2). The interventions were administered orally at least twice daily for one month.

Results: The frequency of pain intensity and mucositis in the two groups was similar on the 14th and 28th days (100% of patients were free from pain and mucositis). Additionally, no significant difference was observed between the two groups in terms of pain intensity and frequency of mucositis on the 7th and 21th days (p > 0.05). On the 7th day, the frequency of grade 1 mucositis was 3.4% in group 1 and 3.3% in group 2, with no statistically significant difference in the severity of mucositis between the two groups (p = 0.981). On the 21th day, the frequency of grade 3 mucositis was 3.4% in group 1 and 0% in group 2; however, there was no statistically significant difference in the severity of mucositis between the two groups on the 21th day (p = 0.305).

Conclusion: Although there was no statistical difference in reducing pain severity, mucositis frequency, or mucositis grade between the two groups, chlorhexidine was found to be more effective and satisfactory in practice. Therefore, considering its lower cost, chlorhexidine is recommended as a cost-effective option for the treatment of oral mucositis.

Keywords: Chlorhexidine, Mouthwash, Mucositis, Neutropenia.

Introduction

In patients undergoing chemotherapy and/or radiotherapy, oral mucositis (OM) is a major acute side effect that impacts the oral cavity (1). The incidence of chemotherapy-induced mucositis is approximately 40% for standard chemotherapy, around 75% for intensive chemotherapy, about 30 to 60% for head and neck radiotherapy, and 90% when radiotherapy is combined with

chemotherapy (2). Importantly, the pain mucositis caused bv oral from radiotherapy and chemoradiotherapy presents a significant negative impact for patients (3). Symptoms can be severe, often resulting in hospitalization and the feeding need for tubes. These severely complications diminish the quality of life for cancer patients and can lead to more serious issues, such as systemic infections. Oral complications like mucositis not only cause intense pain and feeding difficulties but also increase the risk of infection and, in extreme cases, sepsis. Additionally, these complications can elevate treatment costs and necessitate prompting hospital stays, sometimes physicians adjust chemotherapy to dosages. This scenario not only heightens morbidity and mortality rates but also undermines optimal treatment (2, 3). Few pharmacological agents or interventions are effective in alleviating the severity of oral mucositis and the pain associated with radiotherapy (3). The effects of various mouthwashes differ based on their mechanisms of action. Chlorhexidine, a potent antiseptic, combats a wide range of microorganisms, including bacteria and some fungi, and is commonly used to prevent and treat oral infections (4). While chlorhexidine primarily serves as an antiseptic, its impact on pain relief or inflammation is less direct. In contrast, a combination mouthwash containing diphenhydramine, aluminumnystatin, magnesium hydroxide, and lidocaine offers a multifaceted approach. This formulation provides antimicrobial action while also delivering anti-inflammatory, analgesic antifungal, and properties, potentially proving more effective in and symptoms alleviating preventing mucositis (1). Although several combination mouthwash formulations for managing pain from radiotherapy-related oral mucositis have been available for some time (3), there are currently few placebo-controlled randomized trials assessing the efficacy of these preparations, particularly the diphenhvdraminelidocaine-antacid mouthwash. Given the high prevalence of cancer, including acute lymphoblastic leukemia (ALL), in Iran, particularly in Yazd province (5), and considering the use of chemotherapy as the first line of treatment, there is a lack of comprehensive studies on this topic in our country, especially in Yazd province. This study aims to compare the effects of chlorhexidine mouthwash with a combined mouthwash (four drugs) in the prevention and treatment of oral mucositis in neutropenic patients hospitalized in the pediatric oncology department.

Materials and Methods

Design of study and procedure

This study is a double-blind randomized controlled trial (RCT) conducted on children undergoing chemotherapy in the pediatric oncology department of Shahid Sadoughi Hospital, Yazd. The participants included neutropenic patients (<1500 neutrophils per microliter) aged 2 to 18 years receiving chemotherapy. Patients were randomly (using random number table) assigned to two groups of 30: one group received a combined mouthwash containing 25 cc of diphenhydramine, one complete vial of nystatin (100,000 units per milliliter), 25 cc of aluminummagnesium hydroxide, and 4 drops of 2% lidocaine (group 1), while the other group received a 0.2% chlorhexidine mouthwash 2). The interventions (group were administered orally at least twice daily for month. with weekly clinical one examinations conducted by a pediatric oncology specialist who was unaware of the mouthwash type. The mouthwashes in were provided identical coded containers by a nurse who was also unaware of the contents. A designated individual monitored the ethical conduct of the study.

Sampling Procedure

The sample size was determined using the following formula, assuming a 30% difference in the incidence of mucositis between the two groups, with a significance level of 5% and a power of 80%, resulting in a total of 60 participants being assessed.

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

$$n = \frac{\left(Z_{1-\frac{\alpha}{2}} + Z_{1-\beta}\right)^2 \left[P_1(1-P_1) + P_2(1-P_2)\right]}{(P_1 - P_2)^2}$$

Eq. (1)

Inclusion Criteria

1. Patients aged 2 to 18 years.

2. Patients with acute lymphoblastic leukemia (ALL) undergoing chemotherapy with a single protocol (B-cell or T-cell based on the 2022 guidelines).

3. Neutropenia without oral mucositis.

Exclusion Criteria

1. Irregular medication use.

2. Patients who did not return for timely follow-up.

3. Patients experiencing any side effects.

Pain was classified using the Visual Analog Scale (VAS) questionnaire (6). The questions of questionnaire were responded by children.

Degree of Mucositis

According to WHO criteria (7):

• Grade 0: No mucositis.

• **Grade 1:** Mild irritation of the oral mucosa accompanied by pain; no visible lesions present; the patient is able to maintain a normal diet

• Grade 2: Ulcers are present in the oral mucosa; the patient can still swallow pediatric oncology food. Clinical examinations were conducted by a physician unaware of the mouthwash type.

• **Grade 3:** The patient experiences severe sensitivity when swallowing solid food; a liquid diet is essential.

• **Grade 4:** The patient is unable to swallow; complete intravenous or tube feeding is required.

Fig 1 shows consort flowchart of the effect of chlorhexidine mouthwash and combined mouthwash (4 drugs) in the prevention and treatment of oral mucositis in neutropenic patients.

Statistical Analysis

Data were entered into SPSS software version 23. Frequency distributions were reported as frequency and percentage. The Chi-square test was used for statistical analysis, and the independent t-test was employed to compare quantitative data. A p-value of less than 0.05 was considered statistically significant.

Results

The present study was conducted on 59 children with an average age of 6.13 ± 3.6 years. The frequency distribution of these patients in terms of gender showed that 32 (54.2%) were boys and 27 (45.8%) were girls. Additionally, the frequency of side effects was zero in both groups. The mean duration of hospitalization was 4.45 ± 4.71 days in group 1 and 3.50 ± 0.7 days in group 2 (p = 0.761). The comparison of the two groups in terms of pain is shown in Table I. As shown in Table I, the frequency of pain intensity in the two groups was the same on the 14th and 28th days (100% of patients were without pain). Additionally, no significant difference was observed between the two groups in terms of pain intensity on the 7th and 21th days (p > 0.05). The comparison of the two groups in terms of mucositis is shown in Table II. As shown in Table II, the frequency of mucositis in the two groups was the same on the 14th and 28th days (100% of patients were without mucositis). Additionally, no significant difference was observed between the two groups in terms of the frequency of mucositis on the 7th and 21th days (p > 0.05). Table III shows the comparison of the two groups in terms of severity of grade. As the results show, on the 7th day, the frequency of grade 1 mucositis was 3.4% in group 1 and 3.3% in group 2, with no statistically significant difference in the severity of mucositis between the two groups (p = 0.981).

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 Unported License, which permits unrestricted use and redistribution provided that the original author and source are credited.

418

Moreover, the intensity of mucositis was the same in both groups on the 14th and 28th days, with 100% of patients in grade 0. On the 21th day, the frequency of grade 3 mucositis was 3.4% in group 1 and 0%

in group 2; however, there was no statistically significant difference in the severity of mucositis between the two groups on the 21th day (p = 0.305).

Table 1: The comparison of the two groups in terms of pain						
Group	Day	Without pain N (%)	With pain N (%)	Total N (%)	P-value*	
Group 1	7	28 (96.6)	1 (3.4)	29 (100)	0.091	
Group 2	7	29 (96.7)	1 (3.3)	30 (100)	0.981	
Group 1	14	29 (100)	0 (0)	29 (100)		
Group 2	14	30 (100)	0 (0)	30 (100)	•••••	
Group 1	21	28 (96.6)	1 (3.4)	29 (100)	0.305	
Group 2	21	30 (100)	0 (0)	30 (100)	0.305	
Group 1	28	29 (100)	0 (0)	29 (100)		
Group 2	28	30 (100)	0 (0)	30 (100)	•••••	

Combined Mouthwash (group 1), Chlorhexidine mouthwash (group 2) *Chi-Square test

The Type of drug	Day	Without mucositis N (%)	With mucositis N (%)	Total N (%)	P-value
Group 1	7	28	1	29	
Oloup 1	/	(96.6)	(3.4)	(100)	0.981
Group 2	7	29	1	30	
		(96.7)	(3.3)	(100)	
Group 1	14	29	0	29	
		(100)	(0)	(100)	
Group 2	14	30	0	30	
		(100)	(0)	(100)	
Group 1	21	28	1	29	0.305
		(96.6)	(3.4)	(100)	
Group 2	21	30	0	30	
		(100)	(0)	(100)	
Group 1	28	29	0	29	
		(100)	(0)	(100)	
Group 2	28	30	0	30	
		(100)	(0)	(100)	

Table Π : The comparison of the two groups in terms of mucositis

Combined Mouthwash (group 1), Chlorhexidine mouthwash (group 2) *Chi-Square test

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

Comparison of the Effect of Chlorhexidine Mouthwash and Combined Mouthwash (4 drugs) in Preventing and Managing Oral Mucositis in Neutropenic Patients Admitted to the Pediatric Oncology Department: A Randomized Clinical Trial

		1 5	0 1	5	2 3 0 (,	1
The Type of drug		Severity of grade			Total		
	Day	0 N (%)	1 N (%)	2 N (%)	3 N (%)	N (%)	P-value
Group 1	7	28 (96.6)	1 (3.4)	0 (0)	0 (0)	29 (100)	0 981
Group 2	7	29 (96.7)	1 (3.3)	0 (0)	0 (0)	30 (100)	0.901
Group 1	14	29 (100)	0 (0)	0 (0)	0 (0)	29 (100)	
Group 2	14	30 (100)	0 (0)	0 (0)	0 (0)	30 (100)	
Group 1	21	28 (96.6)	0 (0)	0 (0)	1 (3.4)	29 (100)	0 305
Group 2	21	30 (100)	0 (0)	0 (0)	0 (0)	30 (100)	0.305
Group 1	28	29 (100)	0 (0)	0 (0)	0 (0)	29 (100)	_
Group 2	28	30 (100)	0 (0)	0 (0)	0 (0)	30 (100)	

Table III: The comparison of the two groups in terms of severity of grade (mucositis)

Combined Mouthwash (group 1), Chlorhexidine mouthwash (group 2)

*Chi-Square test



Figure 1. Consort flowchat

420

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

Discussion

Oral mucositis is significant a complication of cancer treatment. Various interventions for the prevention and treatment of oral mucositis have been explored. but none have proven completely successful. This study aimed to investigate the effects of chlorhexidine mouthwash and a combination mouthwash (containing four drugs) on the prevention and treatment of oral mucositis in hospitalized neutropenic patients, with no differences noted in their length of stay. Pain frequency assessments in the two groups showed that on the 14th and 28th days, none of the patients in either group reported pain. Additionally, comparisons of pain severity on the 7th and 21th days statistically indicated no significant between the two difference groups. However, on the 21th day, 3.4% of individuals in the four-drug combination (diphenhydramine, group nystatin, aluminum-magnesium hydroxide, and lidocaine) reported pain, while none in the chlorhexidine group experienced pain. Sio et al., studied the effects of a combination mouthwash containing diphenhydramine, aluminum-magnesium lidocaine. hydroxide (an antacid), doxepin (25 mg/5 ml water) on oral mucositis treatment compared to a placebo. They observed that pain scores in both groups four hours after using either mouthwash decreased compared to the control group (3). Torhal et al., examined the efficacy of a mouthwash in reducing discomfort caused by mucositis in chemotherapy patients.

Their mouthwash consisted of three drugs (lidocaine, diphenhydramine, and sodium bicarbonate in normal saline), and patient responses were reported using a selfassessment scale. The results showed that this three-drug mouthwash effectively relieved pain in patients with chemotherapy-induced mucositis, which aligns with our study's findings (8). Limeira et al., investigated and compared chlorhexidine and a combination drug (including nystatin. dexamethasone. diphenhydramine, morphine, lidocaine, B vitamins, and normal saline) against placebo in reducing oral mucositis in a rat model. They observed significant clinical improvements in both intervention groups the compared to placebo group. Furthermore, comparison between the two intervention groups indicated that the combination drug was more effective than chlorhexidine, regarding clinical and histological parameters. In contrast, our study found no statistical difference between the two groups, possibly due to the more extensive components of the combination drug in their study (1). Samiraninezhad et al., studied the effect of mouthwash on pain levels in patients, following them for 14 days. The control group received aluminum-magnesium hydroxide and diphenhydramine, while another group was treated with doxepin, and a third and fourth groups received chitosan nanogel and doxepin / chitosan nano-gel. Results showed that the group treated with doxepin and chitosan nano-gel significantly reduced pain compared to the control group three days post-treatment. Therefore, these results suggest that innovative treatments may be more effective than traditional therapies (9). The comparison of the frequency of mucositis in the two study groups revealed that the frequency of mucositis on the 14th and 28th days was the same. Moreover, no statistically significant difference was seen between the groups on the 7th and 21th days. Dodd et al. conducted a study comparing effects the of three mouthwashes (salt and soda solution, chlorhexidine, and a magic mouthwash Benadryl, containing lidocaine, and Maalox) on patients with mucositis, following them for 12 days. At the end of the 12th day, no statistically significant

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

Comparison of the Effect of Chlorhexidine Mouthwash and Combined Mouthwash (4 drugs) in Preventing and Managing Oral Mucositis in Neutropenic Patients Admitted to the Pediatric Oncology Department: A Randomized Clinical Trial

difference was found between these groups, regarding signs and symptoms. They also recommended the salt and soda solution due to its lower cost (10). Epstein et al. studied the effectiveness of different mouthwashes (chlorhexidine, nystatin, and solution) in preventing saline oral complications in 86 leukemia patients undergoing chemotherapy or bone marrow transplantation. The study found no difference in the incidence of oral sores or mucositis among the treatment groups (11), which aligns with our study's results. The comparison of mucositis grade frequency between the two study groups showed no statistically significant difference between the groups on the 7th and 21th days. Kuk et al. assessed the effectiveness of a magic mouthwash (containing diphenhydramine, dexamethasone, and nystatin) combined with sucralfate, compared to benzydamine hydrochloride, in alleviating the severity of mucositis symptoms induced by cisplatin chemotherapy. No statistically significant difference in mucositis grades was observed between the two groups (12). Therefore, it appears that the combination of magic mouthwash and sucralfate does not outperform benzydamine hydrochloride for the preventive treatment of oral mucositis, aligning with the results of our study. Barker et al. also examined the effectiveness of diphenhydramine syrup and kaolin-pectin in reducing the severity of radiation-induced mucositis. Each group used one of the mouthwashes four times a day, and the results showed no statistically significant difference in mucositis grade (13). These findings are consistent with our study results. Savizadeh et al., conducted a study comparing the efficacy of topical morphine and magic mouthwash in treating oral mucositis, reporting that both morphine and magic mouthwash effectively reduce the severity of cancer treatment-induced oral mucositis, but

morphine was found to be more effective and satisfactory than magic mouthwash (14). Turhal et al. studied the effectiveness including, mouthwash; lidocaine, of diphenhydramine, and sodium bicarbonate saline solution in patients with in chemotherapy-induced oral mucositis and the results showed that the three-drug mouthwash was effective in reducing mucositis severity or symptomatic relief (8). However, this study presented only pre- and post-intervention results and did not compare this intervention with others. Additionally. no side effects were observed in either group in the present study. Another study also examined the side effects of chlorhexidine in treating oral mucositis and found no serious side effects, which aligns with our findings. According to the researchers, the only observed side effects were tooth staining and changes in taste perception, which were not considered significant (15). mouthwash is a multi-drug Magic combination with various components used to reduce the pain and inflammation of oral mucositis, especially in patients undergoing chemotherapy or radiation therapy. These mouthwashes contain different compounds, each with a specific mechanism of action. Diphenhydramine, a first-generation antihistamine, has significant sedative effects and cholinergic blockade. Lidocaine provides temporary pain relief from minor injuries by inhibiting the initiation and transmission of nerve signals, while also decreasing sodium ion permeability in nerve cell membranes (1). Nystatin is crucial in treating fungal infections (16-20), such as candidiasis (1). In cases of oral mucositis, using a nystatin solution is recommended for both prevention and treatment.

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

Conclusion

The result of this study indicated that although there was no statistical difference in reducing pain severity, mucositis frequency, or mucositis grade between the two groups, chlorhexidine proved to be more effective and satisfactory. Moreover, both groups did not experience side effects. Therefore, considering its lower cost, chlorhexidine is recommended as a cost-effective option for the treatment of oral mucositis.

Ethical Considerations

This study was approved by the Ethics Committee of Shahid Sadoughi University (IR.SSU.MEDICINE.REC.1403.036) and the Iranian registry clinical trial (IRCT20180209038673N8).

Acknowledgments

We thank from staffs of Oncology Department of Shahid Sadoughi University. We did not use AI.

Authors' Contributions

MT, RE, conceived and planned the study. M.Y contributed to the sample preparation. M.T contributed to the analysis and interpretation of the results. MT, RE, and, M.Y wrote the primary draft of the manuscript. All the authors had a critical review of the final draft of the manuscript and accepted it.

Funding

None

Conflict of Interest None

References

1. Limeira RR, Ribeiro IL, Bonan PR, da Nóbrega Alves D, dos Santos Ferreira E, da Costa TK, et al. Effect of multidrug solution for the treatment of chemotherapy-induced oral mucositis in vivo. Saudi Dental J 2023; 35(4):378-385.

2. MohammadSoltani G, Hemmati S, Kamali Nejad M, Latifi S A, Musavazadeh S A. Comparison of radiation-induced mucositis with multiple oral Uulcers based on Iranian traditional medicine: A Review Study. Cmja 2019; 9 (3):3748-3759.

3. Sio TT, Le-Rademacher JG, Leenstra JL, Loprinzi CL, Rine G, Curtis A, et al. Effect of doxepin mouthwash or diphenhydramine-lidocaine-antacid mouthwash vs placebo on radiotherapy-

related oral mucositis pain: the alliance A221304 randomized clinical trial. Jama 2019; 321(15): 1481-1490.

4. Steinsapir KD, Woodward JA. Chlorhexidine keratitis: safety of chlorhexidine as a facial antiseptic. Dermatologic Surgery 2017; 43(1):1-6.

5. Hashemi A, Kokab M, Kamalian M, Zarezadeh M, Sheikhpour E, Azod L. The effect of Aloe vera syrup on prevention of fever and neutropenia in children with acute lymphoid leukemia. JJPHO 2020; 10(3): 144-149.

6. Mousavi FS, Golmakani N. Comparison correlation and between visual analogue scale and different dimensions of the short form of McGill pain questionnaire in post-Cesarean Pain Evaluation. Nurs Midwifery J 2018; 16(9): 622-630.

7. Niscola P, Romani C, Cupelli L, Scaramucci L, Tendas A, Dentamaro T, et al. Mucositis in patients with hematologic malignancies: an overview. Haematologica 2007; 92(2):222-231.

8. Turhal NS, Erdal S, Karacay S. Efficacy of treatment to relieve mucositisinduced discomfort. Supportive care in cancer 2000; 8:55-58.

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

Comparison of the Effect of Chlorhexidine Mouthwash and Combined Mouthwash (4 drugs) in Preventing and Managing Oral Mucositis in Neutropenic Patients Admitted to the Pediatric Oncology Department: A Randomized Clinical Trial

9. Samiraninezhad N, Rezaee M, Gholami A, Amanati A, Mardani M. A novel chitosan-based doxepin nanoformulation for chemotherapy-induced oral mucositis: a randomized, doubleblinded, placebo-controlled clinical trial. Inflammopharmacology 2023; 31(5):2411-2420.

10. Dodd MJ, Dibble SL, Miaskowski C, MacPhail L, Greenspan D, Paul SM, et Randomized clinical trial of the al. effectiveness of 3 commonly used mouthwashes to treat chemotherapyinduced mucositis. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology. 2000; 90(1):39-47.

11. Epstein JB, Vickars L, Spinelli J, Reece D. Efficacy of chlorhexidine and nystatin rinses in prevention of oral complications in leukemia and bone marrow transplantation. Oral surg, oral med, oral pathol 1992; 73(6):682-689.

12. Kuk JS, Parpia S, Sagar SM, Tsakiridis T, Kim D, Hodson DI, et al. A randomized phase III trial of magic mouthwash and sucralfate versus benzydamine hydrochloride for prophylaxis of radiation-induced oral mucositis in head and neck cancer. J Clin Oncol 2011; 29(15):5521-5532.

13. Barker G, Loftus L, Cuddy P, Barker B. The effects of sucralfate suspension and diphenhydramine syrup plus kaolin-pectin on radiotherapy-induced mucositis. Oral Surg Oral Med Oral Pathol 2009: 71(3):288-293.

14. Sarvizadeh M, Hemati S, Meidani M, Ashouri M, Roayaei M, Shahsanai A. Morphine mouthwash for the management of oral mucositis in patients with head and neck cancer. Advanced biomed Res 2015; 4(1):44-48.

15. Cardona A, Balouch A, Abdul MM, Sedghizadeh PP, Enciso R. Efficacy of chlorhexidine for the prevention and treatment of oral mucositis in cancer patients: A systematic review with meta-analyses. J Oral Pathol Med 2017; 46(9):680-688.

16. Howell A, Isaacs D, Halliday R, Australasian Study Group for neonatal infections. Oral nystatin prophylaxis and neonatal fungal infections ADC Fetal and Neonatal Edition. 2009; 94(6): F429-33.

17. Aydemir C, Oguz SS, Dizdar EA, Akar M, Sarikabadayi YU, Saygan S, et al. Randomised controlled trial of prophylactic fluconazole versus nystatin for the prevention of fungal colonisation and invasive fungal infection in very low birth weight infants. ADC Fetal & Neonatal Edition 2011; 96(3): F164-168.

18. Sims ME, Yoo Y, You H, Salminen C, Walther FJ. Prophylactic oral nystatin and fungal infections in very-lowbirthweight infants. American J perinatol1988; 5(01):33-36.

19. Blyth CC, Barzi F, Hale K, Isaacs D. Chemoprophylaxis of neonatal fungal infections in very low birthweight infants: efficacy and safety of fluconazole and nystatin. J Pediatrics child health 2012; 48(9): 846-851.

20. Peterson J, Ranganna R. Preventing Fungal Sepsis in Preterm Infants: The Impact of Changing to Prophylactic Nystatin in a Tertiary Neonatal Unit. J Neonatol 2021; 35(2):82-85.

Iran J Ped Hematol Oncol. 2025, Vol 15, No 2, 416-424

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 Unported License, which permits unrestricted use and redistribution provided that the original author and source are credited.

DOI: 10.18502/ijpho.v15i2.18172